

Metric compatibility and determination in complete metric spaces

ARIS DANIILIDIS, TRI MINH LE, DAVID SALAS

Abstract. It was established in [8] that Lipschitz inf-compact functions are uniquely determined by their local slope and critical values. Compactness played a paramount role in this result, ensuring in particular the existence of critical points. We hereby emancipate from this restriction and establish a determination result for merely bounded from below functions, by adding an assumption controlling the asymptotic behavior. This assumption is trivially fulfilled if f is inf-compact. In addition, our result is not only valid for the (De Giorgi) local slope, but also for the main paradigms of average descent operators as well as for the global slope, case in which the asymptotic assumption becomes superfluous. Therefore, the present work extends simultaneously the metric determination results of [8] and [18].

Key words. Determination of a function, Descent modulus, Metric slope, asymptotic criticality.

AMS Subject Classification *Primary* 49J52 ; 58E05 *Secondary* 03F15 ; 30L15

Contents

1	Introduction	1
1.1	Organization of the manuscript	2
1.2	Notation and terminology.	3
2	Descent moduli: state-of-the-art and extended definitions	4
2.1	Determination in compact spaces	5
2.2	A simple noncompact result: the smooth case	7
2.3	General descent paradigms	11
2.4	Metrically compatible descent moduli	12
3	Main result	17
3.1	Comparison lemmas in complete metric spaces	17
3.2	Determination in complete metric spaces	28

1 Introduction

In [8, Theorem 2.4] the authors have shown that in every metric space, the local slope operator contains sufficient information to determine any continuous inf-compact function with finite slope. Indeed, knowledge of the critical values (values of the function on the set of points where the slope is zero) and knowledge of the slope at every point determine uniquely the function. We hereafter refer to this result as *determination result*. The proof makes use of transfinite induction and is based on a cardinality obstruction. Pertinence of the assumptions was also thereby discussed.

In the follow-up work [7] the authors adopted a much more general framework: they introduced

an abstract notion of descent modulus, based on three axioms (see [7, Definition 3.1] or properties (\mathcal{D}_1) – (\mathcal{D}_3) of forthcoming Definition 2.1) and showed that the result of [8] can be emancipated from the metric structure and fit to a mere topological setting, provided a reasonable notion of *steepest descent* (or other meaningful notion of descent, like *average descent*) is coined. Therefore, instead of considering metric spaces, we can work on probability spaces or Markov chains. However, similarly to [8], an underlying compactness assumption was still required in [7]: the functions for which the result applies should (be continuous and) have compact sublevel sets. This was indeed paramount for the proof of the main result of both works.

The aim of the current work is to eliminate the compactness assumption and use instead completeness together with a control on asymptotic behaviour. This renounces full generality, restricting naturally to the framework of (complete) metric spaces.

Very recently, in the same setting of complete metric spaces, Thibault and Zagrodny in [18] were able to obtain a determination result for the *global slope* (we recall this definition in (2.13)). The proof is highly technical and uses the notion of countably orderable families previously introduced in [12]. For a general function, the global slope is a very restrictive notion (controlling also the asymptotic behavior), but for the class of convex functions it coincides with the local slope and the authors were able to obtain the following powerful convex determination result:

- (convex determination) *Two convex continuous and bounded from below functions with the same slope can only differ by a constant.*

The above result was initially established in Hilbert spaces, see [4] (smooth case) and [17] (non-smooth case). It can also be obtained as a corollary of a more general sensitivity result, derived in [6], which states, roughly speaking, that the slope deviation between two convex functions controls the deviation between the functions themselves. A similar determination result was obtained using proximal operators [19]. All these proofs rely heavily on (sub)gradient descent systems, making crucial use of the Hilbertian structure. However, this drawback no longer appears in [18], where the authors, working directly in metric spaces with the global slope, were able to establish the validity of the above convex determination result in Banach spaces.

Coming back to the present work, we enhance the technique developed in [8] to obtain a general determination result in the setting of complete metric spaces. Comparing with [18], the result not only applies for the global slope (where the interest is essentially limited to the convex determination in a Banach space), but also for the local slope (the definition is recalled in (2.12)) as well as for the main paradigms of average descent operator discussed in Section 2.3. As a consequence, the result applies to a large class of functions (for instance, Lipschitz functions in complete metric spaces). This already hints potential applications in Eikonal equations, or more generally, in Hamilton-Jacobi equations whose viscosity solutions admit an alternative description via slopes (see [13], [14], [15] *e.g.*). A further extension is made by formulating the result in terms of an abstract descent modulus in the spirit of [7], but with an extra property (metric compatibility) to reckon with the given metric (see Definition 2.12). From a practical viewpoint, in a given metric space all reasonable descent moduli are metrically compatible (see also discussion in Section 2.4).

1.1 Organization of the manuscript

The manuscript is organized as follows: in Subsection 1.2 we fix notation and terminology, while in Section 2 we revisit from [7] the definition of an abstract descent modulus and readjust it

(*c.f.* Definition 2.1) to encompass extended real-valued functions in a way that the determination result still holds for inf-compact functions which are continuous in their domain.

Subsection 2.1 resumes the State-of-the-art in this (slightly) more general setting, with the extra benefit that the proofs are now significantly simplified. This is possible because Definition 2.1 is defined in a compatible way with respect to function truncation, see proofs of Lemma 2.3 and Theorem 2.4. We then obtain Corollary 2.5 which readily extends [8, Proposition 2.2] and [8, Theorem 2.4].

In Subsection 2.2 we establish an easy noncompact determination result for the case of smooth functions in a Banach space for the natural descent modulus $T[f] = \|\nabla f\|$. The result illustrates perfectly the need of controlling the asymptotic behaviour and at the same time hints towards the right definition of *asymptotically critical sequence* (see Definition 3.1).

In Subsection 2.3 we present the main paradigms of descent in a metric space which are covered by our main result: the (De Giorgi's) local slope, the global slope, the average descent and the diffusion descent. These paradigms are recalled in Subsection 2.4 and treated in a uniform manner by means of the definition-scheme of a *metrically compatible descent modulus* (Definition 2.12).

The main result is presented in Section 3. Controlling the critical values, the asymptotic behavior and the abstract descent at each point leads to Theorem 3.3 (comparison lemma) and Theorem 3.6 (determination result). We recover the determination result of [18] as a corollary, by applying our result for the global slope, which is a particular case of an abstract descent, since in this case every asymptotically critical sequence is infimizing for the function.

1.2 Notation and terminology.

Throughout this work X is a complete metric space, which will be eventually upgraded to a Banach space in Subsection 2.2. Given any function $f : X \rightarrow \mathbb{R}$ and $r \in \mathbb{R}$ we define by

$$[f \leq r] := \{x \in X : f(x) \leq r\},$$

the r -sublevel set. The strict sublevel set $[f < r]$ is defined analogously. We denote the (effective) domain of f by

$$\text{dom } f := \{x \in X : f(x) < +\infty\}.$$

A function f is lower semicontinuous (in short, lsc) if for every $r \in \mathbb{R}$ the sublevel set $[f \leq r]$ is closed. We say that f is proper if it has at least one nonempty sublevel set, or equivalently, if $\text{dom } f \neq \emptyset$. Further, a function f is called *inf-compact* if the sublevel sets $[f \leq r]$ are compact for all $r < \sup f$. Notice that every lower semicontinuous inf-compact function attains its global minimum.

We further denote by $(\mathbb{R} \cup \{+\infty\})^X$ the set of extended real-valued functions on X and by $\mathcal{C}(X)$ the set of continuous real-valued functions on X . We also denote by

$$\begin{aligned} \mathcal{LSC}(X) &:= \{f : X \rightarrow \mathbb{R} \cup \{+\infty\} : f \text{ proper, lsc}\}; \\ \bar{\mathcal{C}}(X) &:= \{f : X \rightarrow \mathbb{R} \cup \{+\infty\} : f \text{ proper, lsc and } f|_{\text{dom } f} \text{ continuous}\}. \end{aligned}$$

A subset \mathcal{F} of $(\mathbb{R} \cup \{+\infty\})^X$ is called a *cone*, if for any $f \in \mathcal{F}$ and $r \geq 0$ we have $rf \in \mathcal{F}$ (with the convention $0 \cdot (+\infty) = 0$). A cone \mathcal{F} which is closed under translation (that is, $f + c \in \mathcal{F}$ for all

$f \in \mathcal{F}$ and $c \in \mathbb{R}$) will be called *translation cone*. In what follows, \mathcal{F} will always be a translation cone of proper functions.

Notice that $\mathcal{C}(X)$ is a vector subspace of $(\mathbb{R} \cup \{+\infty\})^X$ while $\mathcal{LSC}(X), \bar{\mathcal{C}}(X)$ are translation cones. Notice further that $f \in \mathcal{C}(X)$ if and only if $f \in \bar{\mathcal{C}}(X)$ and $\text{dom } f = X$.

For any $a \in \mathbb{R}$ we set $a^+ := \max\{a, 0\}$ (the positive part of the number a).

For a function $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$ we define the operator $\Delta^+ f : \text{dom } f \times X \rightarrow \mathbb{R}$ by

$$\Delta^+ f(x, y) = \begin{cases} \frac{\{f(x) - f(y)\}^+}{d(x, y)}, & \text{if } x \neq y, \\ 0, & \text{if } x = y. \end{cases} \quad (1.1)$$

2 Descent moduli: state-of-the-art and extended definitions

Following the spirit of [7], we call *descent modulus* on a topological space X any operator $T : \mathcal{F} \rightarrow [0, +\infty]^X$ satisfying three natural properties (see (\mathcal{D}_1) – (\mathcal{D}_3) in Definition 2.1 below). The quantity $T[f](x) \in [0, +\infty]$ is conceived as an abstract measurement of descent for the function f at the point x . If $T[f](x) = 0$, then the point x is called *T-critical* (or simply critical). Therefore, the set of *T-critical* points of f coincides with the zeros of the function $T[f]$ and is denoted by

$$\mathcal{Z}_T(f) := \{x \in X : T[f](x) = 0\}. \quad (2.1)$$

A formal definition for proper extended real-valued functions follows:

Definition 2.1 (Descent modulus). *Let $\mathcal{F} \subset (\mathbb{R} \cup \{+\infty\})^X$ be a translation cone. An operator $T : \mathcal{F} \rightarrow [0, +\infty]^X$ is called descent modulus on \mathcal{F} if*

$$\text{dom } T[f] \subset \text{dom } f, \quad \text{for every } f \in \mathcal{F} \quad (\mathcal{D}_0)$$

and the following three conditions hold for every $f, g \in \mathcal{F}$ and $x \in X$:

$$(\mathcal{D}_1) \quad x \in \text{argmin } f \implies x \in \mathcal{Z}_T(f).$$

$$(\mathcal{D}_2) \quad T[f](x) < T[g](x) \implies \exists z \in \text{dom } g : \{f(x) - f(z)\}^+ < \{g(x) - g(z)\}^+.$$

$$(\mathcal{D}_3) \quad \text{If } 0 < T[f](x) < +\infty \text{ and } r > 1, \text{ then } T[f](x) < T[rf](x).$$

Let us have a brief discussion on the properties defining the descent modulus: Property (\mathcal{D}_1) guarantees preservation of global minima, since $\mathcal{Z}_T(f) = \text{argmin } T[f]$. Property (\mathcal{D}_2) can be seen as a monotonicity property on the sublevel set: indeed, if for every $z \in [g \leq g(x)]$ one has $f(x) - f(z) \geq g(x) - g(z)$ (that is, if f has more descent than g in all descent directions of g) then one should necessarily have $T[f](x) \geq T[g](x)$. Therefore, (\mathcal{D}_2) can be restated as follows:

$$\left. \begin{array}{l} \{g(x) - g(z)\}^+ \leq \{f(x) - f(z)\}^+ \\ \text{for all } z \in \text{dom } g \end{array} \right\} \implies T[g](x) \leq T[f](x). \quad (\widetilde{\mathcal{D}}_2)$$

Finally (\mathcal{D}_3) is a scalar monotonicity property, ensuring that if a function f has a nonzero finite descent at x , then the function $(1 + \varepsilon)f$ has an amplified descent at the same point for any $\varepsilon > 0$.

Remark 2.2. (i). Definition 2.1 applies to extended real-valued functions and (\mathcal{D}_0) imposes an infinite descent to all points for which $f(x) = +\infty$. If $\mathcal{F} = \mathcal{C}(X)$, then (\mathcal{D}_0) holds trivially and the above definition of descent modulus coincides with [7, Definition 3.1].

(ii). A straightforward consequence of the above definition is that a descent modulus can be defined only for proper functions. (To see this, given $g \in \mathcal{F}$, consider the function $f \equiv \mathbf{0}$ in \mathcal{F} and apply (\mathcal{D}_2)).

We define the domain $\text{dom } T \subset \mathcal{F}$ of a descent modulus T as follows:

$$\text{dom } T := \{f \in \mathcal{F} : \text{dom } T[f] = \text{dom } f\}, \quad (2.2)$$

that is, $f \in \text{dom } T$ if and only if it has a finite slope at every point in which it has a finite value. If X is a metric space, then $\text{dom } T$ contains the class of Lipschitz continuous functions for every reasonable descent modulus. (The reader can easily verify that this is the case for the main instances of descent moduli of this work: *c.f.* Example 2.10 and Example 2.13.)

2.1 Determination in compact spaces

The determination result established in [7] requires the functions to have compact sublevel sets. The proof was based on a transfinite induction and the conclusion was obtained by contradiction, due to a cardinality obstruction since the induction did not allow point repetitions. In this section, for the sake of completeness, we restate this result in a slightly more general setting: the descent modulus is now considered on extended real-valued (inf-compact) functions. In fact, this new framework, contemplated by the extended Definition 2.1 allows a much simpler proof (namely, the transfinite induction is replaced by a maximum principle), which in the setting of [8], [7] was formally impossible. We present this proof here.

Lemma 2.3 (Strict comparison in compact spaces). *Let X be a compact topological space and T a descent modulus on a translation cone \mathcal{F} containing $\mathcal{LSC}(X)$. Let $f \in \overline{\mathcal{C}}(X)$, $g \in \mathcal{LSC}(X)$ and assume:*

- (i). (*descent domination*) $T[f](x) < T[g](x)$, for every $x \in \text{dom } g \setminus \mathcal{Z}_T(g)$;
- (ii). (*control of criticality*) $f(z) < g(z)$, for every $z \in \mathcal{Z}_T(g)$;

Then, it holds

$$f(x) < g(x), \quad \forall x \in \text{dom } g.$$

Proof. Notice that $\text{dom } g \subset \text{dom } f$, therefore f is continuous on $\text{dom } g$. Let us first assume that g is finite, that is, $\text{dom } g = X$. Then, $f - g$ is (finite and) upper semicontinuous and attains its maximum at some point $x_0 \in X$. It suffices to show that $x_0 \in \mathcal{Z}_T(g)$ (then (ii) applies and $\max(f - g) = (f - g)(x_0) < 0$). If $x_0 \notin \mathcal{Z}_T(g)$, then, $T[g](x_0) > T[f](x_0)$ which yields by hypothesis (\mathcal{D}_2) that there exists $z \in X$ such that $\{f(x_0) - f(z)\}^+ < \{g(x_0) - g(z)\}^+$. In particular, $(f - g)(x_0) < (f - g)(z)$, which is a contradiction.

Let us now consider the case $\text{dom } g \neq X$, that is, g takes the value $+\infty$ at some point. Let $h : X \rightarrow \mathbb{R} \cup \{+\infty\}$ given by

$$h(x) = \begin{cases} (f - g)(x), & \text{if } x \in \text{dom } g, \\ +\infty, & \text{otherwise.} \end{cases}$$

Fix any $a > \inf g$. Since g is lsc, the set $K_a = [g \leq a]$ is nonempty compact and the upper semicontinuous function h attains its maximum there at some point $x_a \in K_a$. If $x_a \notin \mathcal{Z}_T(g)$, then, as before, there exists $z_a \in \text{dom } g$ such that $\{f(x_a) - f(z_a)\}^+ < \{g(x_a) - g(z_a)\}^+$. This yields that $z_a \in K_a$ and $h(x_a) < h(z_a)$, which is a contradiction. Thus, $x_a \in \mathcal{Z}_T(g)$ and h is strictly negative in K_a . Since $\text{dom } g = \bigcup_{a > \inf g} [g \leq a]$, the conclusion follows. \square

The following theorem is the direct extension of the determination theorems of [7], invoking Lemma 2.3 instead of [7, Lemma 3.3].

Theorem 2.4 (Descent determination of extended real-valued functions in compact spaces). *Let X be a compact topological space and T a descent modulus on a translation cone \mathcal{F} containing $\mathcal{LSC}(X)$. Let $f \in \overline{\mathcal{C}}(X)$ and $g \in \mathcal{LSC}(X) \cap \text{dom}(T)$. Then,*

- (a) *If $T[f](x) \leq T[g](x)$, for all $x \in X$ and $f(x) \leq g(x)$, for all $x \in \mathcal{Z}_T(g)$, then $f \leq g$.*
- (b) *If $f, g \in \overline{\mathcal{C}}(X) \cap \text{dom}(T)$, $T[f](x) = T[g](x)$, for all $x \in X$ and $f(x) = g(x)$ for all $x \in \mathcal{Z}_T(g) = \mathcal{Z}_T(f)$, then $f = g$.*

Proof. Since statement (b) is symmetric, it is sufficient to prove (a). Notice that (\mathcal{D}_2) implies that $T[g + c] = T[g]$, for every $c \in \mathbb{R}$ (see also [7, Proposition 3.2(b)]). Therefore, replacing, if necessary, g by $g - \inf g$ and f by $f - \inf g$ we may assume that g is non-negative on X . Now, replacing g by $g_\varepsilon = (1 + \varepsilon)(g + \varepsilon)$, we get that $\text{dom } g_\varepsilon = \text{dom } g$, $T[f](x) < T[g_\varepsilon](x)$ for every $x \in \text{dom } g_\varepsilon$, $\mathcal{Z}_T(g_\varepsilon) \subset \mathcal{Z}_T(g)$ and $f(x) < g_\varepsilon(x)$ for every $x \in \mathcal{Z}_T(g_\varepsilon)$. Thus, $f < g_\varepsilon$ over $\text{dom } g_\varepsilon$. Taking $\varepsilon \rightarrow 0$, we obtain $f \leq g$ on $\text{dom } g$, which readily yields $f \leq g$ on the whole space X . \square

The comparison principles and the determination result of [7] for inf-compact functions can be derived from Lemma 2.3 and Theorem 2.4, as the following corollary shows.

Corollary 2.5. *Let X be a topological space (not necessarily compact) and T a descent modulus on $\mathcal{F} = \overline{\mathcal{C}}(X)$. Let $f, g \in \mathcal{F}$ be bounded from below.*

- (a) *If g is inf-compact, $T[f](x) < T[g](x)$ for all $x \in X \setminus \mathcal{Z}_T(g)$, and $f(x) < g(x)$ for all $x \in \mathcal{Z}_T(g)$, then $f < g$.*
- (b) *If g is inf-compact, $g \in \text{dom}(T)$, $T[f](x) \leq T[g](x)$, for all $x \in X$ and $f(x) \leq g(x)$, for all $x \in \mathcal{Z}_T(g)$, then $f \leq g$.*
- (c) *If f, g are inf-compact, $f, g \in \overline{\mathcal{C}}(X) \cap \text{dom}(T)$, $T[f](x) = T[g](x)$, for all $x \in X$ and $f(x) = g(x)$ for all $x \in \mathcal{Z}_T(g) = \mathcal{Z}_T(f)$, then $f = g$.*

Proof. It is enough to prove (a). The conclusion is trivial if g is constant, since in this case $\mathcal{Z}_T(g) = X$. Therefore we may assume that $\inf g < \sup g$. Fix any $a \in (\inf g, \sup g)$ and set $K_a = [g \leq a]$. Then, K_a is nonempty and compact. We set $\mathcal{F}_a := \{h \in \mathcal{F} : K_a \subset \text{dom } h\}$ and for every $h \in \mathcal{F}$, we define

$$h_a := h + i_{K_a},$$

where i_{K_a} denotes the indicator function of K_a , that is,

$$i_{K_a}(x) := \begin{cases} 0, & x \in K_a \\ +\infty, & x \notin K_a. \end{cases}$$

Notice that $h_a \in \overline{\mathcal{C}(X)} = \mathcal{F}$ and $\mathcal{F}_a \subset \mathcal{F}$, so the operator T is a descent modulus on \mathcal{F}_a . We deduce easily from (\mathcal{D}_2) that for every $h \in \mathcal{F}_a$

$$T[h_a](x) \leq T[h](x), \quad \text{for all } x \in K_a$$

and (since K_a is a nontrivial sublevel set of g)

$$\mathcal{Z}_T(g_a) = \mathcal{Z}_T(g) \cap K_a \quad \text{and} \quad T[g_a](x) = T[g](x), \quad \text{for all } x \in K_a.$$

Then,

$$T[f_a](x) \leq T[f](x) < T[g](x) = T[g_a](x).$$

Moreover, for every $x \in \mathcal{Z}_T(g_a)$ we have $f_a(x) = f(x) < g(x) = g_a(x)$. Applying Lemma 2.3 we deduce that $f_a < g_a$ on K_a , and consequently, $f(x) < g(x)$, for every $x \in K_a$. Since $a \in (\inf g, \sup g)$ is arbitrary, we conclude that $f < g$ on $\text{dom } g \setminus \text{argmax } g$.

If $\text{argmax } g = \emptyset$ or $\sup g = +\infty$, we have $\text{dom } g \setminus \text{argmax } g = \text{dom } g$ and the result follows. Thus, it suffices to consider the case $\sup g < +\infty$ and $\text{argmax } g \neq \emptyset$. Then take $x \in \text{argmax } g$. If $T[g](x) = 0$, then $f(x) < g(x)$ by hypothesis. If not, $T[f](x) < T[g](x)$ and property (\mathcal{D}_2) entails that there exists $z \in \text{dom } g$ such that

$$f(x) - f(z) \leq \{f(x) - f(z)\}^+ < \{g(x) - g(z)\}^+ = g(x) - g(z).$$

Then, $z \notin \text{argmax } g$, entailing that $g(z) > f(z)$. Thus, $f(x) < g(x) + f(z) - g(z) < g(x)$. We conclude that, regardless the value of $T[g](x)$, we always have that $f(x) < g(x)$. Therefore $f < g$ on $\text{dom } g$, and the proof is complete. \square

2.2 A simple noncompact result: the smooth case

Let us consider the setting given by $X = \mathbb{R}^d$, the cone $\mathcal{F} = \mathcal{C}^1(\mathbb{R}^d)$ of continuously differentiable functions, and the descent modulus given by $T[f](x) = \|\nabla f(x)\|$.

In this setting, consider two functions $f, g \in \mathcal{F}$ bounded from below such that

$$\|\nabla f(x)\| \leq \|\nabla g(x)\|, \quad \forall x \in X.$$

Following [17]¹ we compare the functions f and g along the descent curves of g (which is the function with dominating slope). Given x_0 , we consider the curve $\gamma : [0, +\infty) \rightarrow \mathbb{R}^d$ that solves

$$\begin{cases} \dot{\gamma}(t) = -\nabla g(\gamma(t)), & t \geq 0, \\ \gamma(0) = x_0. \end{cases} \quad (2.3)$$

¹A first version of this idea was due to J.-B. Baillon in 2018, see [3].

Then, we can directly write

$$\begin{aligned}
g(x_0) - f(x_0) &= \limsup_{t \rightarrow +\infty} (g - f)(\gamma(t)) - \int_0^t ((g - f) \circ \gamma)'(s) ds \\
&= \limsup_{t \rightarrow +\infty} (g - f)(\gamma(t)) - \int_0^t \left(\langle \nabla g(\gamma(s)), \dot{\gamma}(s) \rangle - \langle \nabla f(\gamma(s)), \dot{\gamma}(s) \rangle \right) ds \\
&= \limsup_{t \rightarrow +\infty} (g - f)(\gamma(t)) + \int_0^t \|\nabla g(\gamma(s))\| |\dot{\gamma}(s)| ds + \int_0^t \langle \nabla f(\gamma(s)), \dot{\gamma}(s) \rangle ds \quad (2.4) \\
&\geq \limsup_{t \rightarrow +\infty} (g - f)(\gamma(t)) + \int_0^t \underbrace{\left(\|\nabla g(\gamma(s))\| - \|\nabla f(\gamma(s))\| \right)}_{\geq 0} |\dot{\gamma}(s)| ds \\
&\geq \limsup_{t \rightarrow +\infty} (g - f)(\gamma(t)) = \lim_{t \rightarrow +\infty} g(\gamma(t)) - \liminf_{t \rightarrow +\infty} f(\gamma(t)).
\end{aligned}$$

Consequently, if there exists $\bar{x} \in \omega\text{-lim } \gamma$ (ω -limit point of γ), then $\bar{x} \in Z := \mathcal{Z}_T(g)$. In such a case, if $g|_Z \geq f|_Z$ (which is the boundary condition of Theorem 2.4 (a)), we would conclude that $g(x_0) \geq f(x_0)$. However, since the space is noncompact, $\omega\text{-lim } \gamma$ might be empty and we need to include the boundary condition

$$\liminf_{t \rightarrow +\infty} f(\gamma(t)) \leq \lim_{t \rightarrow +\infty} g(\gamma(t)). \quad (2.5)$$

The above condition should be imposed only for steepest descent curves $\gamma : [0, +\infty) \rightarrow \mathbb{R}^d$ of g without ω -limits and not for those for which $\omega\text{-lim } \gamma \neq \emptyset$, since this latter case is already captured by the comparison condition on the critical set $\mathcal{Z}_T(g)$ (ω -limits of the gradient descent curve are automatically critical points for g).

The drawback of the boundary condition (2.5) is that it depends on ∇g , via (2.3), rather than on the descent modulus $T[g] = \|\nabla g\|$. To overcome this difficulty and obtain a boundary condition that is independent of ∇g we introduce the following definition:

Definition 2.6 (Asymptotically critical path, smooth version). *We say that a differentiable curve $\tilde{\gamma} : [0, +\infty) \rightarrow \mathbb{R}^d$ is an asymptotically T -critical path for the function g (with $T[g] = \|\nabla g\|$) if*

$$\|\tilde{\gamma}'\| = 1, \quad \omega\text{-lim } \tilde{\gamma} = \emptyset \quad \text{and} \quad \int_0^{+\infty} T[g](\tilde{\gamma}(s)) ds < +\infty. \quad (2.6)$$

The above definition encompasses the following key elements for the class of continuously differentiable bounded from below functions:

- (1) If $\tilde{\gamma}$ is an asymptotically critical path for g , then $\lim_{s \rightarrow \infty} T[g](\tilde{\gamma}(s)) = 0$.
- (2) Every steepest descent curve without ω -limits yields, upon reparametrization, an asymptotically critical path.
- (3) If two continuously differentiable functions have the same slope, then they have the same critical points and the same asymptotically critical paths.

The last assertion is obvious from Definition 2.6, while the first is a straightforward consequence of the integrability condition in (2.6). Concerning the second assertion, let $\gamma : [0, +\infty) \rightarrow \mathbb{R}^d$ be

a descent curve of g without ω -limits. Then, γ has infinite length and so does $\tilde{\gamma}$, its arc-length parametrization, defined by

$$\tilde{\gamma}(s) := \gamma(\sigma^{-1}(s)) \quad \text{where} \quad s = \sigma(t) := \int_0^t \|\dot{\gamma}(t)\| dt. \quad (2.7)$$

Then, $\omega\text{-lim } \tilde{\gamma} = \emptyset$ and $\|\tilde{\gamma}'(s)\| = 1$. Performing a change of variables we deduce:

$$g(\gamma(0)) - \inf g \geq - \int_0^{+\infty} (g \circ \gamma)'(t) dt = \int_0^{+\infty} \|\nabla g(\gamma(t))\| \cdot \|\dot{\gamma}(t)\| dt = \int_0^{+\infty} \|\nabla g(\tilde{\gamma}(s))\| ds, \quad (2.8)$$

which yields (2.6).

With the above in mind, we establish the following noncompact determination result.

Theorem 2.7. *Let $f, g : \mathbb{R}^d \rightarrow \mathbb{R}$ be two continuously differentiable functions which are bounded from below. Assume that*

- (i) $\|\nabla f(x)\| = \|\nabla g(x)\|$ for every $x \in \mathbb{R}^d$;
- (ii) $f(z) = g(z)$ for every $z \in \mathcal{Z}_T(f) = \mathcal{Z}_T(g)$.
- (iii) $\liminf_{s \rightarrow +\infty} f(\tilde{\gamma}(s)) = \liminf_{s \rightarrow +\infty} g(\tilde{\gamma}(s))$, for each (common) asymptotically critical path $\tilde{\gamma}$.

Then, $f = g$.

Proof. Take $x_0 \in \mathbb{R}^d$ and γ be a steepest descent curve of g emanating from x_0 . Then, we can divide our analysis in two cases:

Case 1: $\omega\text{-lim } \gamma$ is nonempty. Choose $\bar{z} \in \omega\text{-lim } \gamma$. Since $\lim_{t \rightarrow \infty} \|\nabla g(\gamma(t))\| = 0$, continuity of ∇g entails that $\bar{z} \in \mathcal{Z}_T(g)$. Moreover, continuity of g and f yield that $\lim_{t \rightarrow \infty} g(\gamma(t)) = g(\bar{z})$ and $\lim_{t \rightarrow \infty} f(\gamma(t)) = f(\bar{z})$. Then, (2.4) yields that

$$g(x_0) - f(x_0) \geq g(\bar{z}) - f(\bar{z}) = 0.$$

Case 2: $\omega\text{-lim } \gamma$ is empty.

Then, since γ is the steepest descent curve of g emanating from x_0 , we get that

$$\int_0^{+\infty} \|\nabla g(\gamma(t))\| \cdot \|\dot{\gamma}(t)\| dt = - \int_0^{+\infty} (g \circ \gamma)'(t) dt = g(x_0) - \lim_{t \rightarrow +\infty} g(\gamma(t)) \leq g(x_0) - \inf g < +\infty.$$

Thus, the curve $\tilde{\gamma}$ given in (2.7) is an asymptotically critical path and consequently assumption (iii) applies. We deduce from (2.4) that

$$g(x_0) - f(x_0) \geq \lim_{t \rightarrow +\infty} g(\gamma(t)) - \liminf_{t \rightarrow +\infty} f(\gamma(t)) = \lim_{s \rightarrow +\infty} g(\tilde{\gamma}(s)) - \liminf_{s \rightarrow +\infty} f(\tilde{\gamma}(s)) = 0.$$

In either case we get $g(x_0) \geq f(x_0)$ and (since x_0 is arbitrary) deduce $g \geq f$. Exchanging the roles of g and f , we obtain the desired result. \square

Remark 2.8 (importance of integrability condition). It is tempting to simplify Definition 2.6 and replace the integrability condition in (2.6) on the path $\gamma : [0, +\infty) \rightarrow \mathbb{R}^d$ by its consequence

$$\lim_{s \rightarrow +\infty} T[g](\gamma(s)) = 0 \quad (2.9)$$

(see (1) after Definition 2.6 above), possibly together with the requirement that the curve is unbounded with no ω -limits (to avoid redundancy with critical points). The following example illustrates that this would lead to an undesirably large class of asymptotically critical curves:

Indeed, set $\mathcal{U} := \mathbb{R} \times (0, +\infty)$ and consider the convex function

$$\begin{cases} g : \mathcal{U} \rightarrow \mathbb{R} \\ g(x, y) = \frac{x^2}{y}. \end{cases} \quad (2.10)$$

Then, $\text{Im}(g) = [0, +\infty)$ and $c = 0$ is the unique critical value of g (in particular, every point of the level set $[g = 0] := \{0\} \times (0, +\infty)$ is critical!). Since g is a convex and $\inf g = 0$, the result of [4] applies and g is determined by its slope $\|\nabla g\|$ and its minimum value $\inf g = 0$. Let us now focus on assumption (iii) of Theorem 2.7 (control on asymptotic critical paths). For a convex function, the integrability condition (2.6) yields $g(\gamma(s)) \rightarrow \inf g = 0$. Therefore, (iii) is not an additional requirement (it is already contained in assumption (ii)) and Theorem 2.7 generalizes the convex determination result mentioned in the introduction.




Figure 1: Function $g(x, y) = x^2/y$. In blue, the plane $z = 1$. In black, the curve $\gamma(t)$ for $c = 1$. In dashed red, the curve $(\gamma(t), g(\gamma(t))) = (\gamma(t), c)$ for $c = 1$. Plane XY plotted in 1:6, starting at point (1,1). Plane XZ plotted in 1:1, starting at point (1,0).

Omitting the integrability condition in Definition 2.6 would have led to a completely different situation: for every $c > 0$, the level set

$$[g = c] := \{(x, y) \in \mathcal{U} : x^2 = cy\} = \{(\sqrt{ct}, t^2) : t \neq 0\}$$

contains an unbounded curve $\gamma(t) = (\sqrt{ct}, t^2)$, $t > 0$, without ω -limits, satisfying

$$g(\gamma(t)) = c \quad \text{and} \quad \nabla g(\gamma(t)) = \left(\frac{2\sqrt{c}}{t}, \frac{c}{t^2} \right) \xrightarrow{t \rightarrow \infty} 0,$$

that is, every $c > 0$ would have been an asymptotic critical value of g (see Figure 1 for an illustration of the case $c = 1$). Therefore, assumption (iii) of Theorem 2.7 becomes very restrictive leading to an essentially useless statement. \diamond

In what follows, we will move to metric spaces and establish determination results for general classes of functions. The lack of derivatives (and norms) is addressed by the metric slope, which we consider under an abstract unified framework encompassing several other paradigms of descent operators. An additional difficulty is to control the asymptotic behaviour of the functions: asymptotically critical paths will be replaced by asymptotically critical sequences $\{z_n\}_n$ with $T[g](z_n) \rightarrow 0$ as $n \rightarrow \infty$ and the integrability condition (2.6) by the summability condition

$$\sum_{n \geq 1} T[g](z_n) d(z_n, z_{n+1}) < +\infty. \quad (2.11)$$

Indeed, since $\tilde{\gamma}$ is parameterized by arc-length, setting $z_n := \tilde{\gamma}(s_n)$, for all $n \geq 1$, we deduce

$$s_{n+1} - s_n \geq d(\tilde{\gamma}(s_{n+1}), \tilde{\gamma}(s_n))$$

and (2.11) follows directly from the discretization of (2.6). This together with the fact that the sequence has no accumulation points eventually leads to Definition 3.1.

2.3 General descent paradigms

Our goal is to derive a nonsmooth version of Theorem 2.7 for functions defined in a complete metric space (X, d) . As already mentioned, this requires a suitable extension of the notion of asymptotically critical paths in order to impose a boundary condition in the lines of condition (iii) of Theorem 2.7. Moreover, we aim to obtain a statement that generalizes (and recovers) the determination results of [8] (local slope, inf-compact case) and [18] (global slope, complete metric case) and apply to the main paradigms of descent moduli studied in [7] and quoted below:

1. *The local (metric) slope* ([10], [1] e.g.): For any $f \in \mathcal{LSC}(X)$ the local (metric) slope is given by:

$$s[f](x) = s_f(x) := \begin{cases} \limsup_{y \rightarrow x} \Delta^+ f(x, y), & \text{if } x \in \text{dom } f \\ +\infty, & \text{otherwise.} \end{cases} \quad (2.12)$$

2. *The global slope*: Similarly, for any $f \in \mathcal{LSC}(X)$ the global slope is given by

$$\mathcal{G}[f](x) := \begin{cases} \sup_{y \in X} \Delta^+ f(x, y), & \text{if } x \in \text{dom } f \\ +\infty, & \text{otherwise.} \end{cases} \quad (2.13)$$

3. *The average descent modulus*: Let μ be a probability measure over X , and let \mathcal{F} be the μ -measurable extended-valued functions on X . The average descent modulus is then given by

$$\mathcal{M}[f](x) = \begin{cases} \int_X \Delta_f^+(x, y) \mu(dy) = \int_X \{f(x) - f(y)\}^+ \left(\frac{1}{d(x, y)} \mu(dy) \right), & \text{if } x \in \text{dom } f, \\ +\infty, & \text{otherwise.} \end{cases} \quad (2.14)$$

This is an oriented nonlocal operator determined by the family of measures $\{\mu_x\}_x$ with

$$\mu_x(dy) = \frac{1}{d(x,y)}\mu(dy), \text{ if } y \neq x \quad (\text{under the convention } \frac{0}{0} = 0).$$

(see [7, Definition 4.14]).

4. *The diffusion descent modulus:* Let \mathcal{F} be the μ -measurable extended-valued functions on X , and suppose now that $\mu(A) > 0$ for every open set A of X . Then, we define the diffusion descent modulus \mathcal{D} over \mathcal{F} given by

$$\mathcal{D}[f](x) = \begin{cases} \limsup_{\varepsilon \rightarrow 0^+} \frac{1}{\mu(B(x, \varepsilon))} \int_{B(x, \varepsilon)} \Delta_f^+(x, y) \mu(dy), & \text{if } x \in \text{dom } f, \\ +\infty, & \text{otherwise.} \end{cases} \quad (2.15)$$

This is the oriented 1-dispersion operator for measure μ (see [7, Definition 4.2]).

All four notions described above fit the definition of descent modulus in the extended sense of Definition 2.1. The proofs are mild adaptations of [7]. In all cases, Lipschitz continuous functions are contained in the domain of each of the aforementioned descent moduli.

2.4 Metrically compatible descent moduli

The definition of a descent modulus (*cf.* Definition 2.1) does not require prior assumptions on the space X . In particular, X does not need to be metric (neither topological) space, although the aforementioned determination result in [7, Theorem 3.5] will eventually require topology, to formulate the assumptions of continuity and compactness. This being said, this work is inscribed in the framework of a complete metric space (X, d) . In order to obtain determination results in this setting and ensure an efficient use of completeness property, we need to impose some (metric) compatibility condition to the considered descent modulus reckoning with completeness of X . The condition should encompass the four paradigms of Subsection 2.3. The first natural attempt for such condition leads to the following definition.

Definition 2.9 (strong metric compatibility). *We say that a descent modulus $T : \mathcal{F} \rightarrow [0, +\infty]^X$ is strongly metrically compatible, if for some strictly increasing continuous function $\theta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\theta(0) = 0$ and $\lim_{t \rightarrow \infty} \theta(t) = +\infty$ and for every $f, g \in \text{dom}(T)$, $x \in \text{dom } g$ and $\delta > 0$ it holds:*

$$T[f](x) < \delta < T[g](x) \quad \implies \quad \exists z \in \text{dom } g : \frac{\{f(x) - f(z)\}^+}{d(x, z)} < \theta(\delta) < \frac{\{g(x) - g(z)\}^+}{d(x, z)} \quad (\widehat{\mathcal{D}}_2)$$

The idea behind the above definition is to guarantee that whenever $T[f](x) < \delta < T[g](x)$, there exists a point z ensuring on the one hand more descent for g than for f and on the other hand enough descent for g relative to the distance $d(x, z)$ (up to a factor $\theta(\delta)^{-1}$).

It is straightforward to see that $(\widehat{\mathcal{D}}_2)$ yields (\mathcal{D}_2) . Another important remark is that the function θ is invertible, and its inverse $\theta^{-1} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ verify the same properties as θ , that is, θ is a strictly increasing continuous function with $\theta^{-1}(0) = 0$ and $\lim_{s \rightarrow \infty} \theta^{-1}(s) = +\infty$.

The above definition encompasses the *sup*-type paradigms of metric descent modulus.

Example 2.10 (steepest descent operators). (i). (local slope) Let us recall from (2.12) the definition of the local slope. It has been shown in [7, Proposition 3.7] that the “local slope” operator

$$s : (\mathbb{R} \cup \{+\infty\})^X \rightarrow [0, +\infty]^X,$$

defined by $s[f] := s_f$ is a descent modulus on the linear subspace $\mathcal{C}(X)$, that is, it satisfies properties (\mathcal{D}_1) – (\mathcal{D}_3) of Definition 2.1. Since (\mathcal{D}_0) is also verified, it follows easily that $s[\cdot]$ is a descent modulus on the translation cone $\mathcal{F} := \mathcal{LSC}(X)$ as well. Moreover, let $f, g \in \mathcal{LSC}(X)$ and $x \in \text{dom } g$ be such that $s_f(x) < \delta < s_g(x)$. Then, (2.12) yields that for some $\sigma > 0$ sufficiently small, we have:

$$\sup_{y \in B(x, \sigma)} \Delta_f^+(x, y) < \delta < \sup_{y \in B(x, \sigma)} \Delta_g^+(x, y).$$

In particular, we can choose $z \in B(x, \sigma)$ such that $\Delta_g^+(x, z) > \delta$, and obtain

$$\frac{\{f(x) - f(z)\}^+}{d(x, z)} < \delta < \frac{\{g(x) - g(z)\}^+}{d(x, z)}.$$

Thus $(\widehat{\mathcal{D}}_2)$ holds for $\theta(\delta) = \delta$.

(ii). (global slope) The global slope of a function $f \in \mathcal{LSC}(X)$ at a point $x \in X$ is defined in (2.13) as follows:

$$\mathcal{G}_f(x) := \sup_{y \in X} \Delta_f^+(x, y) = \sup_{y \in \text{dom } f} \Delta_f^+(x, y)$$

It is straightforward to see that the “global slope” operator $\mathcal{G} : \mathcal{LSC}(X) \rightarrow [0, +\infty]^X$ is strongly metrically compatible, satisfying $(\widehat{\mathcal{D}}_2)$ with $\theta(\delta) = \delta$. \diamond

Notice that if a descent modulus T is strongly metrically compatible, then $\hat{T} := \phi \circ T$ remains strongly metrically compatible, for every strictly increasing continuous function with $\phi(0) = 0$ and $\lim_{t \rightarrow +\infty} \phi(t) = +\infty$. However, unfortunately, Definition 2.9 is quite restrictive and fails to cope with some important *average*-type paradigms, as reveals the following example:

Example 2.11 (Average descent fails strong metric compatibility). Set $X = [0, 1]$ endowed with the distance $d(t, s) = |t - s|$ and the usual Lebesgue measure. We show that for any function $\theta : [0, +\infty) \rightarrow [0, +\infty)$ given as in Definition 2.9 and any $\delta > 0$, there exist functions f and g for which $(\widehat{\mathcal{D}}_2)$ fails for the average descent modulus \mathcal{M} (see Subsection 2.3). Indeed, fix $\theta(\cdot)$, $\delta > 0$ and pick any positive number $\varepsilon < \min \left\{ \frac{\delta}{2}, \frac{\theta(\delta)}{6} \right\}$, then set

$$t_0 = \min \left\{ \frac{1}{2}, \frac{\delta - \varepsilon}{2\theta(\delta)} \right\}, \quad t_1 = t_0 + \frac{3\varepsilon}{2\theta(\delta)}, \quad t_2 = t_0 + \frac{3\varepsilon}{\theta(\delta)}.$$

We can easily see that $\delta - \varepsilon > 0$ and $t_2 < 1$. We now define $h > 0$ in a way that

$$\frac{1}{2} (h - \theta(\delta)) t_0 + \theta(\delta) t_0 = \delta - \varepsilon,$$

that is, $h = \frac{2(\delta-\varepsilon)}{t_0} - \theta(\delta)$. By the definition of t_0 , it is easy to check that $h > 3\theta(\delta)$. Let us define functions $\psi, \phi : X \rightarrow [0, +\infty)$ as follows:

$$\psi(t) = \begin{cases} h - \frac{h-\theta(\delta)}{t_0} t, & \text{if } t \in [0, t_0) \\ \theta(\delta) - \frac{\theta(\delta)^2}{3\varepsilon}(t - t_0), & \text{if } t \in [t_0, t_2], \\ 0, & \text{if } t \in [t_2, 1]. \end{cases} \quad \text{and} \quad \phi(t) = \begin{cases} h - \frac{h-\theta(\delta)}{t_0} t, & \text{if } t \in [0, t_0), \\ \theta(\delta) - \frac{2\theta(\delta)^2}{3\varepsilon}(t - t_0), & \text{if } t \in [t_0, t_1], \\ 0, & \text{if } t \in [t_1, 1]. \end{cases}$$

It is not hard to realize (see Figure 2) that

$$\int_X \psi(t) dt = \delta + \frac{\varepsilon}{2} > \delta > \delta - \frac{\varepsilon}{4} = \int_X \phi(t) dt.$$




Figure 2: *Left:* Regions below the functions ψ and ϕ . *Right:* The functions $g(t) = -t\psi(t)$ and $f(t) = -t\phi(t)$ coincide on $[0, t_0]$ while for $t \geq t_0$ we have $f \geq g$.

Now, we can define the functions $g, f : X \rightarrow \mathbb{R}$ given by

$$g(t) = -t\psi(t) \quad \text{and} \quad f(t) = -t\phi(t).$$

Then ψ and ϕ coincide with the functions $\Delta^+g(0, \cdot)$ and $\Delta^+f(0, \cdot)$. Indeed, since g and f attain their global maximum at $t = 0$, we can directly write

$$\Delta^+g(0, t) = \frac{g(0) - g(t)}{t} = \psi(t) \quad \text{and} \quad \Delta^+f(0, t) = \frac{f(0) - f(t)}{t} = \phi(t).$$

Thus, $(\widehat{\mathcal{D}}_2)$ fails at 0, since $\mathcal{M}[f](0) = \delta - \frac{\varepsilon}{4} < \delta < \delta + \frac{\varepsilon}{2} < \mathcal{M}[g](0)$ but there is no $t \in X$ such that $\phi(t) < \theta(\delta) < \psi(t)$. Even worse, there is no $t \in X$ such that $\phi(t) < \psi(t)$ (more descent for g than for f) and $\theta(\delta) < \psi(t)$ (enough descent for g) simultaneously. \diamond

The previous example reveals that average-type descents fail Definition 2.9 (strong metric compatibility). To overcome this difficulty, we need to relax this definition as follows:

Definition 2.12 (metric compatibility). A descent modulus $T : \mathcal{F} \rightarrow [0, +\infty]^X$ is said to be metrically compatible, if for every $\rho > 0$, there exists a strictly increasing continuous function $\theta_\rho : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\theta_\rho(0) = 0$ and $\lim_{t \rightarrow +\infty} \theta_\rho(t) = +\infty$, such that for every $f, g \in \text{dom}(T)$, $x \in \text{dom } g$ and $\delta > 0$ it holds:

$$T[f](x) < \delta < T[g](x) \implies \exists z \in \text{dom } g : \begin{cases} \{f(x) - f(z)\}^+ < (1 + \rho)\{g(x) - g(z)\}^+ \\ \text{and} \\ \theta_\rho(\delta) d(x, z) < g(x) - g(z) \end{cases} \quad (C)$$

The above definition is a trade-off between the needs of the proof of Theorem 3.3 and a common scheme that incorporates all main paradigms of Subsection 2.3. The difference with Definition 2.9 is that given $\rho > 0$, if $T[f](x) < \delta < T[g](x)$, we can find a point z (depending on ρ) ensuring more descent for g than for f up to a factor $(1 + \rho)$ and sufficient descent for g relative to the distance $d(x, z)$, up to a factor $\theta_\rho(\delta)^{-1}$ (depending again on ρ). In this sense, for a given tolerance $\rho > 0$, condition (C) is a trade-off between these requirements.

Average descent operators are operators for which the descent of f at a point x is obtained by integrating the quotient $\Delta_f^+(x, y)$ with respect to some probability measure on X . This category of operators are now metrically compatible with respect to this relaxed definition:

Example 2.13 (average descent moduli). Let μ be a probability measure on the metric space (X, d) and consider the translation cone

$$\mathcal{F} := \{f : X \rightarrow \mathbb{R} \cup \{+\infty\} : f \text{ is proper and } \mu\text{-measurable}\}.$$

(i) Consider the operator \mathcal{M} defined in (2.14). Let us show that \mathcal{M} is a metrically compatible descent modulus. Indeed, fix $\rho > 0$ and assume that for some $f, g \in \text{dom}(\mathcal{M})$, $x \in \text{dom } g$ and $\delta > 0$ we have $\mathcal{M}[f](x) < \delta < \mathcal{M}[g](x)$. Then, by definition, we have that

$$\int_X (1 + \rho) \Delta_g^+(x, y) \mu(dy) > \int_X (\Delta_f^+(x, y) + \rho \delta) \mu(dy) = \int_X \Delta_f^+(x, y) d\mu + \rho \delta,$$

which yields that there exists $z \in \text{dom } g$ such that $(1 + \rho)\Delta_g^+(x, z) > \Delta_f^+(x, z) + \rho\delta$. The above inequality yields

$$(1 + \rho)\{g(x) - g(z)\}^+ > \{f(x) - f(z)\}^+ \quad \text{and} \quad \frac{g(x) - g(z)}{d(x, z)} > \underbrace{\left(\frac{\rho}{1 + \rho}\right)}_{\theta_\rho(\delta)} \delta.$$

Thus, (C) is verified for $\theta_\rho(\delta) := \frac{\rho}{1 + \rho} \delta$.

(ii) Suppose now that $\mu(A) > 0$ for every open set A of X and consider the oriented 1-dispersion operator for the measure μ , given by (2.15). Then, \mathcal{D} is a metrically compatible descent modulus. Indeed, let $f, g \in \text{dom}(\mathcal{D})$, $x \in \text{dom } g$ and $\delta \in \mathbb{R}$ such that $\mathcal{D}[f](x) < \delta < \mathcal{D}[g](x)$. Then, for $\varepsilon > 0$ sufficiently small we have

$$\frac{1}{\mu(B(x, \varepsilon))} \int_{B(x, \varepsilon)} \Delta_f^+(x, y) \mu(dy) < \delta < \frac{1}{\mu(B(x, \varepsilon))} \int_{B(x, \varepsilon)} \Delta_g^+(x, y) \mu(dy).$$

Then, the conclusion follows by noting that $\nu(dy) = \frac{1}{\mu(B(x, \varepsilon))} \mu(dy)$ is a probability measure over the metric space $B(x, \varepsilon)$, and therefore we can proceed as in the previous example (i). \diamond

Based on these examples-schemes, we can significantly enlarge the class of metrically compatible descent moduli as follows: for any strictly increasing, continuous function $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\phi(0) = 0$ and $\lim_{t \rightarrow +\infty} \phi(t) = +\infty$, we can replace $\Delta_f^+(x, y)$ (in definitions (2.12), (2.13), (2.14) and (2.15)) by

$$\widetilde{\Delta}_f^+(x, y) := \phi^{-1}(\Delta_f^+(x, y))$$

and obtain new classes.

Let us now give an example of a descent modulus which is not metric compatible.

Example 2.14. Take $X = \mathbb{N}$ endowed with the distance function d given by $d(m, n) = |m - n|$. Consider the operator T given by

$$T[f](k) = \sup_{m \in \mathbb{N}} \{f(k) - f(m)\}^+, \quad \text{for all } k \in \mathbb{N}.$$

Clearly, T is a descent modulus (it coincides with the global slope for the distance d_0 given by $d_0(k, m) = 1$, if $k \neq m$, and $d_0(k, m) = 0$, if $k = m$). Consider the function f_n , $n \in \mathbb{N}$, given by

$$f_n(m) = \begin{cases} 1, & \text{if } m \neq n, \\ 0, & \text{if } m = n. \end{cases}$$

Observe that for each $\delta \in (0, 1)$ and each $n \geq 2$, we have that $T[f_n](1) = 1 > \delta > 0 = T[f_1](1)$. However, regardless the tolerance $\rho > 0$, the only choice for $m \in [f_n < f_n(1)]$ is $m = n$ and so

$$(1 + \rho) \Delta_{f_n}^+(1, n) = (1 + \rho) \frac{f_n(1) - f_n(n)}{d(1, n)} = \frac{1 + \rho}{n - 1} > 0 = \frac{\{f_1(1) - f_1(n)\}^+}{d(1, n)} = \Delta_{f_1}^+(1, n).$$

If T were metrically compatible, there would exist a continuous, strictly increasing function $\theta_\rho : [0, +\infty) \rightarrow [0, +\infty)$, with $\theta_\rho(0) = 0$ and $\theta_\rho(\delta) < \frac{f_n(1) - f_n(n)}{d(n, 1)} = \frac{1}{n-1}$ for all $\delta \in (0, 1)$ and all $n \geq 2$, which is a contradiction. Therefore T is not metrically compatible for the metric d . Notice however, that it is so for the discrete metric d_0 . \diamond

Remark 2.15. We recall from [7, Section 3.4] that there exist slope-like operators, as the weak slope ([5], [9] e.g.) or the limiting slope ([11]) that are not descent moduli, since they fail property (\mathcal{D}_2) (monotonicity).

We finish the section with two stability properties that we will need in the sequel. The first one stems from [7, Proposition 3.2(b)], whose proof is based on the monotonicity property (\mathcal{D}_2) of the descent moduli.

Proposition 2.16 (translation-invariance). *Let $T : \mathcal{F} \rightarrow [0, +\infty]^X$ be a descent modulus. Then for any $f \in \mathcal{F}$ and $c \in \mathbb{R}$, we have*

$$T[f] = T[f + c].$$

The second property provides another way of constructing new metrically compatible descent moduli, similar to the remark after Example 2.13.

Proposition 2.17 (composition with increasing functional). *Let $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a strictly increasing continuous function with $\phi(0) = 0$ and $\lim_{t \rightarrow +\infty} \phi(t) = +\infty$. Let*

$$T : \mathcal{F} \rightarrow [0, +\infty]^X$$

be a descent modulus. One has that

$$T \text{ is metrically compatible} \iff \phi \circ T \text{ is metrically compatible}, \quad (2.16)$$

where $\phi \circ T$ is the descent modulus given by $(\phi \circ T)[f](x) = \phi(T[f](x))$, with $\phi(+\infty) = +\infty$.

Proof. The fact that $\phi \circ T$ is a descent modulus was established in [7, Proposition 3.9] for real-valued functions. The proof can be easily adapted to the present setting of extended real-valued functions. Concerning metric compatibility, since ϕ^{-1} is also strictly increasing and continuous, with $\phi^{-1}(0) = 0$ and $\lim_{s \rightarrow +\infty} \phi^{-1}(s) = +\infty$, it is sufficient to establish only one implication. To this end, let us assume that T is metrically compatible with $\{\theta_\rho\}_{\rho>0}$ given as in Definition 2.12. Let $f, g \in \text{dom}(\phi T)$, $x \in \text{dom} g$ and $\delta > 0$ such that $(\phi T)[f](x) < \delta < (\phi T)[g](x)$. Fix $\rho > 0$. It is straightforward to see that $f, g \in \text{dom}(T)$ and $T[f](x) < \phi^{-1}(\delta) < T[g](x)$, therefore for some $z \in \text{dom} g$ we have

$$\{f(x) - f(z)\}^+ < (1 + \rho)\{g(x) - g(z)\}^+ \quad \text{and} \quad \theta_\rho(\phi^{-1}(\delta)) < \frac{g(x) - g(z)}{d(x, z)}.$$

Since $\tilde{\theta}_\rho := \theta_\rho \circ \phi^{-1}$ is continuous and strictly increasing with $\tilde{\theta}_\rho(0) = 0$ and $\lim_{t \rightarrow +\infty} \tilde{\theta}_\rho(t) = +\infty$, the conclusion follows. \square

3 Main result

This section contains our main result: we establish a comparison principle, in the lines of Lemma 2.3, for metrically compatible descent moduli (*cf.* Definition 2.12) in a complete, but not necessarily compact, metric space. This result will eventually lead to our determination result (Theorem 3.6). Absence of compactness imposes an assumption on the asymptotic behaviour for which, as already discussed in Subsection 2.2, the choice of the notion of asymptotic criticality is paramount. This latter not only consists of saying that the descent moduli vanish at infinity, but also requires two additional restrictions: absence of accumulation points and a summability condition (Definition 3.1). The price to pay is that the scheme of proof becomes more involved, but as a reward, we are able to obtain a statement that generalizes all previous results (*c.f.* Theorem 3.6). Both the local slope determination of [8] for inf-compact functions and the global slope determination [18] are now recovered by our final statement.

3.1 Comparison lemmas in complete metric spaces

In this section, we want to establish a comparison principle, in the lines of Lemma 2.3, for metrically compatible descent moduli (*cf.* Definition 2.12) in a complete (but not necessarily compact) metric space. Absence of compactness assumption leads inevitably to impose control on the asymptotic behavior. To do so, we need a precise notion of asymptotic T -criticality, which is given in the following definition:

Definition 3.1 (asymptotic critical sequences). *A sequence $\{z_n\}_{n \geq 1} \subset X \setminus \mathcal{Z}_T(g)$ is called T -asymptotically critical for a function $g \in \mathcal{F}$ (in short, $T[g]$ -critical) if it has no converging subsequence and*

$$\sum_{n=0}^{+\infty} T[g](z_n) d(z_n, z_{n+1}) < +\infty. \quad (3.1)$$

We denote by $\mathcal{AZ}_T(g)$ the set of asymptotically critical sequences for g .

Remark 3.2 (justification of terminology). Any sequence $\{z_n\}_{n \geq 1}$ satisfying $\liminf_{n \rightarrow \infty} T[g](z_n) > 0$ and (3.1) is necessarily Cauchy: indeed, assume that for some $\delta > 0$ and $N \geq 1$ we have $T[g](z_n) \geq \delta$, for all $n \geq N$. Then for every $\varepsilon > 0$, we take $n_0 \geq N$ such that

$$\sum_{i \geq N} T[g](z_i) d(z_i, z_{i+1}) < \varepsilon \delta,$$

and for every $m > n \geq n_0$ we deduce

$$d(z_n, z_m) \leq \sum_{i=n}^{m-1} d(z_i, z_{i+1}) \leq \frac{1}{\delta} \sum_{i=n}^{m-1} T[g](z_i) d(z_i, z_{i+1}) < \varepsilon.$$

Therefore, absence of convergent subsequences in Definition 3.1 ensures that in a complete metric space, every $T[g]$ -critical sequence $\{z_n\}_n$ should satisfy

$$\liminf_{n \rightarrow \infty} T[g](z_n) = 0. \quad (3.2)$$

Another important feature of this notion is that it becomes vacuous under compactness.

The requirement for the convergence of the series in (3.1) restricts significantly the class of critical sequences. This was motivated by the discussion in Subsection 2.2 and can be seen as a discrete version of the integrability condition in (2.6), which in turn was inspired by the steepest descent gradient system. A similar condition was also employed in the fundamental lemma [18, Lemma 4.2] used to derive the determination result in [18]. Our next result (Theorem 3.3) extends the metric determination result of [18], since it holds for any metrically compatible descent modulus (and not only for the global slope). Indeed, in case of the global slope we have (see forthcoming Lemma 3.7):

- If $T[g] = \mathcal{G}[g]$ (global slope) and $\{z_n\}_n$ is a $T[g]$ -critical sequence, then $g(z_n) \xrightarrow[n \rightarrow \infty]{} \inf g$.

We are now ready to establish the main comparison lemma.

Theorem 3.3 (comparison lemma). *Let (X, d) a complete metric space and $T : \mathcal{F} \rightarrow [0, +\infty]^X$ a metrically compatible descent modulus. Let $f, g \in \text{dom}(T)$ be two bounded from below functions with $g \in \mathcal{LSC}(X)$ and $f \in \bar{\mathcal{C}}(X)$. Let us assume that:*

- (i). (descent domination) $T[f](x) < T[g](x)$, for every $x \in X \setminus \mathcal{Z}_T(g)$;
- (ii). (control of criticality) $f(z) \leq g(z)$, for every $z \in \mathcal{Z}_T(g)$;
- (iii). For every $\{z_n\}_n \in \mathcal{AZ}_T(g)$ it holds:

$$\liminf_{n \rightarrow +\infty} f(z_n) \leq \liminf_{n \rightarrow +\infty} g(z_n). \quad (3.3)$$

Then, it holds

$$f(x) \leq g(x), \quad \text{for every } x \in \text{dom } g.$$

Proof. We deduce from (i) and (\mathcal{D}_0) of Definition 2.1 that $\text{dom } g \subset \text{dom } f$.

Let us fix $\rho > 0$. Replacing f and g by $f - \inf g + 1$ and $g - \inf g + 1$ if needed, we may assume that $g > 0$ and consequently $(1 + \rho)g > g > 0$. In what follows, we prove $f < (1 + \rho)g$. Once this is done, since ρ is arbitrarily small, we deduce $f \leq g$.

To this end, let $\{\theta_{\rho'}\}_{\rho' > 0}$ be a family of continuous strictly increasing functions associated to the descent modulus T (c.f. Definition 2.12). By Proposition 2.16 the operator $\widehat{T} := \theta_{\rho} \circ T$ is a metrically compatible descent modulus, which preserves T -critical points (i.e. $\mathcal{Z}_{\widehat{T}}(g) = \mathcal{Z}_T(g)$) and $T[g]$ -critical sequences (i.e. a sequence is $T[g]$ -critical if and only if it is $\widehat{T}[g]$ -critical). Furthermore, assumption (i) continues to hold for \widehat{T} and (C) is now satisfied for $\theta_{\rho}(\delta) = \delta$, $\delta > 0$. Therefore, by replacing T by \widehat{T} if necessary, we may assume that for the value $\rho > 0$ (that we fixed in the beginning) condition (C) holds for the identity function $\theta_{\rho}(\delta) = \delta$.

With all the above in mind, let us define, for every $x \in \text{dom } g$, the quantity

$$\delta(x) := \frac{1}{2}T[f](x) + \frac{1}{2}T[g](x). \quad (3.4)$$

Notice that since $f, g \in \text{dom } T$, we have $\delta(x) < +\infty$. Moreover, if $x \notin \mathcal{Z}_T(g)$ (that is, x is not T -critical for g), then (i) yields $\delta(x) > 0$ and

$$T[f](x) < \delta(x) < T[g](x) < 2\delta(x). \quad (3.5)$$

We now assume, towards a contradiction, that there exists $x_0 \in \text{dom } g$ such that

$$f(x_0) \geq (1 + \rho)g(x_0).$$

In what follows, using successively assumption (i) (descent domination) and condition (C) of Definition 2.12, we are going to construct a sequence of points $\{x_n\}_{n \geq 1}$ failing the conclusion of our theorem, where additionally g is strictly decreasing.

Basic iteration scheme (classical induction). Thanks to assumption (ii), we have $x_0 \notin \mathcal{Z}_T(g)$ and we deduce from (i) that (3.5) holds for $x = x_0$. Using (C) for $\theta(\delta) = \delta$ and $\delta = \delta(x_0)$ we obtain $x_1 \in [g < g(x_0)]$ such that

$$0 < f(x_0) - (1 + \rho)g(x_0) \leq f(x_1) - (1 + \rho)g(x_1) \quad \text{and} \quad \delta(x_0)d(x_0, x_1) \leq g(x_0) - g(x_1). \quad (3.6)$$

Therefore we obtain:

$$g(x_0) > g(x_1), \quad f(x_1) > (1 + \rho)g(x_1) \quad \text{and} \quad x_1 \notin \mathcal{Z}_T(g).$$

It is quite clear that the above procedure can be repeated as many times as we wish, producing a sequence $\{x_n\}_{n \geq 1}$ in $X \setminus \mathcal{Z}_T(g)$ such that

$$\text{the sequence } \{(f - (1 + \rho)g)(x_n)\}_{n \geq 0} \text{ is nondecreasing and positive} \quad (3.7)$$

and

$$0 < \delta(x_n)d(x_n, x_{n+1}) < g(x_n) - g(x_{n+1}), \quad \text{for every } n \geq 0. \quad (3.8)$$

The telescopic series obtained by summing up the above inequalities for all $n \geq 0$, together with (3.5) and the fact that the sequence $\{g(x_n)\}_{n \geq 0}$ is strictly decreasing and bounded from below, yield that

$$\frac{1}{2} \sum_{n=0}^{\infty} T[g](x_n) d(x_n, x_{n+1}) < \sum_{n=0}^{\infty} \delta(x_n) d(x_n, x_{n+1}) \leq g(x_0) - \inf g < +\infty. \quad (3.9)$$

Assumption (iii) together with (3.7) ensure that the sequence $\{x_n\}_{n \geq 0}$ cannot be $T[g]$ -critical, therefore we deduce from the above inequality and Definition 3.1 that $\{x_n\}_{n \geq 0}$ has accumulation points as $n \rightarrow \infty$.

First limit ordinal. Let us denote by ω the first infinite ordinal and by $\omega^+ \equiv \omega + 1$ its successor. We first consider the case where

$$\liminf_{n \rightarrow \infty} T[g](x_n) \geq \delta > 0. \quad (3.10)$$

It follows from (3.9) that the sequence $\{x_n\}_{n \geq 1}$ is Cauchy (see Remark 3.2), therefore it converges to some point $\bar{x} \in X$. Setting $x_\omega := \bar{x}$ we deduce easily from (3.8) that

$$g(x_n) > g(\bar{x}), \quad \text{for every } n \geq 0. \quad (3.11)$$

Similarly, we deduce from (3.7) that

$$(f - (1 + \rho)g)(\bar{x}) > 0, \quad (3.12)$$

therefore, $x_\omega \in X \setminus \mathcal{Z}_T(g)$, $\delta(x_\omega) > 0$ and the basic iteration scheme can be pursued from x_ω to define $x_{\omega+1}$, $x_{\omega+2}$ etc.

We now focus on the case

$$\liminf_{n \rightarrow \infty} T[g](x_n) = 0. \quad (3.13)$$

Then, x_ω should be selected inside the set of accumulation points of the sequence $\{x_n\}_{n \geq 0}$ (as we have seen, this set is nonempty, but if (3.13) holds, it might not be a singleton). Although any accumulation point \bar{x} satisfies inequalities (3.11) and (3.12), we cannot assign x_ω randomly among the accumulation points, but instead, we need to select it in an adequate way (for reasons that relate to forthcoming property (P3) required in our forthcoming transfinite induction). To this end, let us set $k_0 = 0$ and define inductively:

$$k_{n+1} := \min \{m \geq k_n : T[g](x_m) < T[g](x_{k_n})\}. \quad (3.14)$$

With this construction, $\{T[g](x_{k_n})\}_{n \in \mathbb{N}}$ is strictly decreasing and $T[g](x_{k_n}) \rightarrow 0$. Since for all $\ell \in [k_n, k_{n+1}) \cap \mathbb{N}$ we have $T[g](x_\ell) \geq T[g](x_{k_n})$ we deduce easily from the triangular inequality that:

$$T[g](x_{k_n}) d(x_{k_n}, x_{k_{n+1}}) \leq \sum_{\ell=k_n}^{k_{n+1}-1} T[g](x_\ell) d(x_\ell, x_{\ell+1})$$

yielding that

$$\sum_{n=0}^{+\infty} T[g](x_{k_n}) d(x_{k_n}, x_{k_{n+1}}) \leq \sigma(\omega) := \sum_{n=0}^{\infty} T[g](x_n) d(x_n, x_{n+1}) < +\infty. \quad (3.15)$$

Therefore, the obtained subsequence $\{x_{k_n}\}_{n \geq 1}$ should also have accumulation points (it cannot be $T[g]$ -critical, thanks to (iii) and (3.7)) and we define x_ω to be any accumulation point \bar{x} of $\{x_{k_n}\}_{n \geq 1}$.

Idea of the proof. Let us outline the main idea of the proof: so far we have defined $\{x_n\}_{n < \omega^+} \equiv \{x_n\}_{n < \omega} \cup \{x_\omega\}$ in $X \setminus \mathcal{Z}_T(g)$ such that $\{g(x_\alpha)\}_{\alpha < \omega^+}$ is strictly decreasing (in terms of ordinals). Let us denote by Ω the first uncountable ordinal, that is,

$$\Omega = \{ \lambda : \lambda \text{ countable ordinal} \}.$$

Our objective is to extend $\{x_n\}_{n < \omega^+}$ to all countable ordinals and come up with a generalized sequence $\{x_\lambda\}_{\lambda < \Omega}$ in $X \setminus \mathcal{Z}_T(g)$ such that $\{g(x_\lambda)\}_{\lambda < \Omega}$ is strictly decreasing. Then for every $\lambda < \Omega$ we would have $g(x_\lambda) - g(x_{\lambda^+}) > 0$ (where λ^+ denotes the successor of λ) and since Ω is uncountable we would deduce:

$$g(x_0) - \inf g \geq g(x_0) - \inf_{\lambda < \Omega} g(x_\lambda) \geq \sum_{\lambda < \Omega} [g(x_\lambda) - g(x_{\lambda^+})] = +\infty. \quad (3.16)$$

The above clearly contradicts the fact that the function g is bounded and proves the result. This construction is naturally based on transfinite induction, where we should (also) deal with ordinals of limit type (ie. $\lambda = \sup\{\alpha : \alpha < \lambda\}$). In this case, x_λ should be defined among the accumulation points of $\{x_\alpha\}_{\alpha < \lambda}$ so that we can deduce $x_\lambda \notin \mathcal{Z}_T(g)$ and guarantee the strict descent of g . However, in absence of compactness, we need an additional argument to ensure that the set accumulation points is nonempty whenever

$$\liminf_{\alpha < \lambda} T[g](x_\alpha) = 0 \quad (\text{else, the limit } \lim_{\alpha \nearrow \lambda} x_\alpha \text{ exists!})$$

In this critical situation, we need to construct a sequence $\{x_{\alpha_n}\}_{n \geq 1}$ with $\alpha_n \nearrow \lambda$ (out of the generalized sequence $\{x_\alpha\}_{\alpha < \lambda}$) satisfying

$$\sum_{n=0}^{+\infty} T[g](x_{\alpha_n}) d(x_{\alpha_n}, x_{\alpha_{n+1}}) < +\infty,$$

use assumption (iii) to deduce that $\{x_{\alpha_n}\}_{n \geq 1}$ cannot be $T[g]$ -critical, evoke Definition 3.1 to conclude that it has accumulation points and eventually select x_λ among them. (Notice that although the set Ω is uncountable—which is crucial for the contradiction in (3.16) above—all of its elements $\lambda < \Omega$ are countable ordinals and consequently, we can always obtain cofinal sequences.) This being said, in order to effectively realize the aforementioned critical step, we need to verify properties (P1)–(P3) below at every step of the forthcoming transfinite induction.

Properties (P1)–(P3): For every ordinal $\lambda \in [\omega, \Omega)$, we are going to construct a generalized sequence $\{x_\alpha\}_{\alpha < \lambda}$ satisfying the following properties:

(P1) For every $0 \leq \alpha < \alpha^+ < \lambda$:

$$0 < \delta(x_\alpha) d(x_\alpha, x_{\alpha^+}) < g(x_\alpha) - g(x_{\alpha^+}). \quad (3.17)$$

Notice that (3.17) yields in particular that

$$\sigma(\lambda) := \sum_{0 \leq \alpha < \lambda} T[g](x_\alpha) d(x_\alpha, x_{\alpha+}) < +\infty. \quad (3.18)$$

(P2) For every $0 \leq \alpha_1 < \alpha_2 < \lambda$:

$$0 \leq (f - (1 + \rho)g)(x_{\alpha_1}) \leq (f - (1 + \rho)g)(x_{\alpha_2}) \quad (3.19)$$

and

$$d(x_{\alpha_1}, x_{\alpha_2}) \leq \sum_{\alpha_1 \leq \alpha < \alpha_2} d(x_\alpha, x_{\alpha+}). \quad (3.20)$$

(P3) For every $\varepsilon > 0$ and ordinals $0 \leq \alpha_0 < \xi < \lambda$, there exists a finite sequence of ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_N < \alpha_{N+1} := \xi$ such that

$$\sum_{n=0}^N T[g](x_{\alpha_n}) d(x_{\alpha_n}, x_{\alpha_{n+1}}) < \sigma(\xi) - \sigma(\alpha_0) + \varepsilon. \quad (3.21)$$

Construction (via transfinite induction). We use transfinite induction as follows: assuming $\{x_\alpha\}_{\alpha < \lambda} \subset X \setminus \mathcal{Z}_T(g)$ is well-defined and satisfies (P1)–(P3), we define $x_\lambda \in X \setminus \mathcal{Z}_T(g)$ in a way that the extended generalized sequence $\{x_\alpha\}_{\alpha < \lambda+} \equiv \{x_\alpha\}_{\alpha \leq \lambda}$ still satisfies the same properties. In case of a successor ordinal $\lambda = \beta^+$, since $x_\beta \notin \mathcal{Z}_T(g)$, defining $x_\lambda \equiv x_{\beta+}$ by means of (C) of Definition 2.12 automatically guarantees that (P1)–(P3) continue to hold for $\{x_\alpha\}_{\alpha < \lambda+}$ (see details below). If λ is a limit-ordinal and x_λ is an accumulation point of the generalized sequence $\{x_\alpha\}_{\alpha < \lambda}$, that is,

$$x_\lambda \in \bigcap_{\alpha < \lambda} \overline{\{x_{\alpha'} : \alpha' \geq \alpha\}} \quad (\text{equivalently, } x_\lambda = \lim_{n \rightarrow \infty} x_{\alpha_n}, \text{ for some } \alpha_n \nearrow \lambda) \quad (3.22)$$

then (P1) is automatically fulfilled by the induction step (since no new successor ordinal is added) and (P2) follows by passing to the limit. Therefore, the main technical difficulty is to show that $\{x_\alpha\}_{\alpha < \lambda}$ has accumulation points and to define x_λ (among these accumulation points) in a way that (P3) holds. Let us now proceed to a rigorous construction:

Initialization. We have already defined $\{x_\alpha\}_{\alpha < \omega+} = \{x_n\}_{n \geq 0}$ satisfying (P1)–(P2). Let us prove that (P3) also holds. Fix $\varepsilon > 0$ and consider the case $\alpha_0 = 0$ and $\xi = \omega$ (every other value of $\alpha_0 \in \mathbb{N}$ follows replacing $\{x_n\}_{n \geq 0}$ by $\{x_n\}_{n \geq \alpha_0}$). We consider three cases:

Case 1: $\liminf_{n \rightarrow \infty} T[g](x_n) = \delta \in (0, +\infty)$. Since (3.9) holds, in view of Remark 3.2, the sequence $\{x_n\}_{n \geq 1}$ converges and $x_\omega = \lim_{n \rightarrow \omega} x_n$. Taking $n_0 \geq 0$ such that

$$d(x_n, x_\omega) < \frac{\varepsilon}{\delta + 1}, \quad \text{for all } n \geq n_0,$$

and choosing $N \geq n_0$ such that $T[g](x_N) < \delta + 1$, we readily obtain

$$\underbrace{\sum_{n=0}^{N-1} T[g](x_n) d(x_n, x_{n+1})}_{< \sigma(\omega)} + T[g](x_N) d(x_N, x_\omega) < \sigma(\omega) + \varepsilon.$$

The result follows for $\alpha_n := n$, $n \in \{0, 1, \dots, N\}$ and $\alpha_{N+1} := \omega$.

Case 2: $\liminf_{n \rightarrow \infty} T[g](x_n) = +\infty$. Similarly to the previous case, the sequence $\{x_n\}_{n \geq 1}$ converges and $x_\omega = \lim_{n \rightarrow \omega} x_n$. Let $N \geq 0$ be such that

$$T[g](x_N) = \min_{n \geq 0} \{T[g](x_n)\} \quad (\text{the minimum is attained since } T[g](x_n) \rightarrow +\infty).$$

Then, $T[g](x_n) \geq T[g](x_N)$ for all $n \geq N$ which in view of (3.20) (for $\alpha_1 = N$ and $\alpha_2 = \omega$) yields

$$T[g](x_N) d(x_N, x_\omega) \leq T[g](x_N) \sum_{n=N}^{+\infty} d(x_n, x_{n+1}) < \sum_{n=N}^{+\infty} T[g](x_n) d(x_n, x_{n+1}).$$

Consequently,

$$\sum_{n=0}^{N-1} T[g](x_n) d(x_n, x_{n+1}) + T[g](x_N) d(x_N, x_\omega) \leq \sum_{n=0}^{+\infty} T[g](x_n) d(x_n, x_{n+1}) := \sigma(\omega).$$

Case 3: $\liminf_{n \rightarrow \infty} T[g](x_n) = 0$. In this case x_ω is defined among the accumulation points of the sequence $\{x_{k_n}\}_{n \geq 1}$ constructed in (3.14). Given $\varepsilon > 0$ we chose $n_0 \in \mathbb{N}$ in a way that

$$d(x_{k_{n_0}}, x_\omega) < \frac{\varepsilon}{T[g](x_{k_{n_0}})}.$$

In view of (3.15), we deduce that

$$\sum_{n=0}^{n_0-1} T[g](x_{k_n}) d(x_{k_n}, x_{k_{n+1}}) + T[g](x_{k_{n_0}}) d(x_{k_{n_0}}, x_\omega) < \sigma(\omega) + \varepsilon,$$

and the result follows for $N = k_{n_0}$ and $\alpha_n := k_n$, $n \in \{0, 1, \dots, N\}$.

Successor ordinal. Assume $\lambda = \beta^+ > \omega$ is a successor ordinal and $\{x_\alpha\}_{\alpha < \lambda} \equiv \{x_\alpha\}_{\alpha \leq \beta}$ is well-defined and satisfies (P1)–(P3). Then from (3.19) and (ii) we deduce that $x_\beta \notin \mathcal{Z}_T(g)$ and $\delta(x_\beta) > 0$. Using (C) as before, we obtain $x_{\beta^+} \in [g < g(x_\beta)]$ such that

$$0 < f(x_\beta) - (1 + \rho)g(x_\beta) \leq f(x_{\beta^+}) - (1 + \rho)g(x_{\beta^+}) \quad \text{and} \quad \delta(x_\beta)d(x_\beta, x_{\beta^+}) \leq g(x_\beta) - g(x_{\beta^+}).$$

Notice that (3.20), (3.21) follow easily from the induction step and the triangular inequality. Therefore $\{x_\alpha\}_{\alpha < \lambda^+} \equiv \{x_\alpha\}_{\alpha \leq \lambda}$ also satisfies (P1)–(P3).

Ordinal of limit type. Let us now assume that $\lambda \in (\omega, \Omega)$ is a limit ordinal and $\{x_\alpha\}_{\alpha < \lambda}$ is defined and satisfies (P1)–(P3).

Case I. We first focus on the case where

$$\liminf_{\alpha < \lambda} T[g](x_\alpha) \geq \delta > 0. \tag{3.23}$$

We deduce that there exists $\xi_0 < \lambda$ such that $T[g](x_\alpha) \geq \delta/2$ for all $\xi_0 \leq \alpha < \lambda$. Thus, in an analogous way as in Remark 3.2, we get from (3.18) that

$$\sum_{\xi_0 \leq \alpha < \lambda} d(x_\alpha, x_{\alpha^+}) \leq \frac{2}{\delta} \sum_{\xi_0 \leq \alpha < \lambda} T[g](x_\alpha) d(x_\alpha, x_{\alpha^+}) < +\infty$$

and consequently the generalized sequence $\{x_\alpha\}_{\alpha < \lambda}$ converges as $\alpha \nearrow \lambda$. We set

$$x_\lambda := \lim_{\alpha \rightarrow \lambda} x_\alpha$$

and obtain readily that the generalized sequence $\{x_\alpha\}_{\alpha < \lambda^+}$ still satisfies (P1)–(P2). It remains to check that (P3) holds for $\varepsilon > 0$ and $0 \leq \alpha_0 < \lambda < \lambda^+$. To this end, we need to consider separately two different cases depending on whether (3.23) is finite or not.

– *Subcase I₁*. Assume that $\liminf_{\alpha < \lambda} T[g](x_\alpha) = \delta \in (0, +\infty)$. Then, we fix $\alpha_* \geq \alpha_0$ such that for all $\alpha \in (\alpha_*, \lambda)$ we have

$$d(x_\alpha, x_\lambda) < \frac{\varepsilon}{2(\delta + 1)}.$$

Pick any $\hat{\alpha} \in (\alpha_*, \lambda)$ such that $T[g](x_{\hat{\alpha}}) < \delta + 1$. We deduce directly that

$$T[g](x_{\hat{\alpha}}) d(x_{\hat{\alpha}}, x_\lambda) < \frac{\varepsilon}{2}.$$

Applying (P3) for $0 \leq \alpha_0 < \hat{\alpha} < \lambda$ (and $\tilde{\varepsilon} = \varepsilon/2$) we obtain a finite strictly increasing sequence $\{\alpha_n\}_{n=0}^N$ satisfying

$$\sum_{n=0}^{N-1} T[g](x_{\alpha_n}) d(x_{\alpha_n}, x_{\alpha_{n+1}}) + T[g](x_{\alpha_N}) d(x_{\alpha_N}, x_{\hat{\alpha}}) < \sigma(\hat{\alpha}) - \sigma(\alpha_0) + \frac{\varepsilon}{2}.$$

The result follows by concatenation.

– *Subcase I₂*. Assume that $\liminf_{\alpha < \lambda} T[g](x_\alpha) = \lim_{\alpha < \lambda} T[g](x_\alpha) = +\infty$. Then, there exists $\alpha_* \geq \alpha_0$ such that $T[g](x_\alpha) \geq 1$, for all $\alpha \in (\alpha_*, \lambda)$. We deduce from (3.18) that

$$\Delta := \sum_{\alpha_* \leq \alpha < \lambda} d(x_\alpha, x_{\alpha^+}) \leq \sum_{\alpha_* \leq \alpha < \lambda} T[g](x_\alpha) d(x_\alpha, x_{\alpha^+}) < +\infty.$$

Set

$$\mu_* := \inf_{\alpha_* \leq \alpha < \lambda} \{T[g](x_\alpha)\} \geq 1.$$

In contrast to Case 2 of Initialization, the above infimum might not be attained if λ is limit of ordinals of limit type (that is, $\lambda = \sup\{\xi < \lambda : \xi \text{ limit-ordinal}\}$). However, we can choose $\hat{\alpha} \in (\alpha_*, \lambda)$ such that

$$T[g](x_{\hat{\alpha}}) < \mu_* + \frac{\varepsilon}{2\Delta}.$$

We deduce from (3.20) (for $\alpha_1 = \hat{\alpha}$ and $\alpha_2 = \lambda$) that

$$\begin{aligned} T[g](x_{\hat{\alpha}}) d(x_{\hat{\alpha}}, x_\lambda) &\leq T[g](x_{\hat{\alpha}}) \sum_{\hat{\alpha} \leq \alpha < \lambda} d(x_\alpha, x_{\alpha^+}) < \sum_{\hat{\alpha} \leq \alpha < \lambda} \left(\mu_* + \frac{\varepsilon}{2\Delta} \right) d(x_\alpha, x_{\alpha^+}) \\ &< \underbrace{\sum_{\hat{\alpha} \leq \alpha < \lambda} T[g](x_\alpha) d(x_\alpha, x_{\alpha^+})}_{:= \sigma(\lambda) - \sigma(\hat{\alpha})} + \frac{\varepsilon}{2}. \end{aligned}$$

We conclude as before by concatenating $\{\hat{\alpha}, x_\lambda\}$ with the finite sequence obtained by applying (P3) (which by the induction step is assumed to hold for the generalized sequence $\{x_\alpha\}_{\alpha < \lambda}$) for the

choice $\tilde{\varepsilon} = \varepsilon/2$ and $0 \leq \alpha_0 < \hat{\alpha} < \lambda$.

Case II. It remains to deal with the case

$$\liminf_{\alpha < \lambda} T[g](x_\alpha) = 0.$$

Our objective is to show that

$$\mathcal{C} = \bigcap_{\alpha < \lambda} \overline{\{x_{\alpha'} : \alpha' \geq \alpha\}} \neq \emptyset$$

and define x_λ in \mathcal{C} in a way that $\{x_\alpha\}_{\alpha < \lambda^+}$ satisfies (P3). (We recall that if $x_\lambda \in \mathcal{C}$ then (P1)–(P2) are automatically satisfied.) To this end, fix $\varepsilon > 0$ and $0 \leq \alpha_0 < \lambda < \lambda^+$. Let

$$\{\varepsilon_n\}_{n \geq 0} \subset (0, \varepsilon) \quad \text{such that} \quad \sum_{n=0}^{+\infty} \varepsilon_n = \frac{\varepsilon}{2}.$$

Let further $\gamma_n \nearrow \lambda$ and define inductively a sequence of ordinals $\{\xi_n\}_{n \geq 0}$ as follows:

$$\xi_0 = \alpha_0 \quad \text{and} \quad \xi_{n+1} := \min \left\{ \alpha \geq \max\{\xi_n, \gamma_n\} : T[g](x_\alpha) \leq \frac{T[g](x_{\xi_n})}{2} \right\}.$$

The above definition guarantees that $\xi_n \nearrow \lambda$ and $T[g](x_{\xi_n}) \searrow 0$ (that is, the sequence of descent moduli converges to 0 decreasingly). For every $n \geq 0$, thanks to our induction assumption, we can apply property (P3) for $\varepsilon_n > 0$ and $0 \leq \xi_n < \xi_{n+1} < \lambda^+$ to obtain $N_n \geq 1$ and a finite sequence $\xi_n := \alpha_0^n < \alpha_1^n < \dots < \alpha_{N_n+1}^n := \xi_{n+1}$ such that

$$\sum_{i=0}^{N_n} T[g](x_{\alpha_i^n}) d(x_{\alpha_i^n}, x_{\alpha_{i+1}^n}) < \sigma(\xi_{n+1}) - \sigma(\xi_n) + \varepsilon_n.$$

Concatenating the above finite sequences $\{\alpha_i^n : i \in \{0, \dots, N_n\}\}$, for $n \geq 0$, we obtain a strictly increasing sequence

$$\alpha_0 := \alpha_0^0 < \alpha_1^0 < \dots < \alpha_{N_0+1}^0 := \xi_1 := \alpha_0^1 < \dots < \alpha_{N_1+1}^1 := \xi_2 < \dots < \dots \quad (3.24)$$

which converges to λ and satisfies

$$\sum_{n=0}^{+\infty} \sum_{i=0}^{N_n} T[g](x_{\alpha_i^n}) d(x_{\alpha_i^n}, x_{\alpha_{i+1}^n}) < \sigma(\lambda) - \sigma(\alpha_0) + \frac{\varepsilon}{2}.$$

Renaming (3.24) to $\{\beta_n\}_{n \geq 0}$, we have $\beta_0 \equiv \alpha_0$, $\beta_n \nearrow \lambda$

$$\sum_{n=0}^{+\infty} T[g](x_{\beta_n}) d(x_{\beta_n}, x_{\beta_{n+1}}) < \sigma(\lambda) - \sigma(\alpha_0) + \frac{\varepsilon}{2} \quad \text{and} \quad \liminf_{n \rightarrow \infty} T[g](x_{\beta_n}) = 0.$$

Acting as in (3.14), we set $k_0 = \beta_0 \equiv \alpha_0$ and

$$k_{n+1} := \min \{\beta_m \geq k_n : T[g](x_{\beta_m}) < T[g](x_{k_n})\}.$$

Since for all $\ell \in [k_n, k_{n+1}) \cap \{\beta_m\}_{m \in \mathbb{N}}$ we have $T[g](x_{\beta_\ell}) \geq T[g](x_{\beta_{k_n}})$, we deduce:

$$\sum_{n=0}^{+\infty} T[g](x_{k_n}) d(x_{k_n}, x_{k_{n+1}}) \leq \sum_{n=0}^{+\infty} \sum_{\ell=k_n}^{k_{n+1}-1} T[g](x_\ell) d(x_\ell, x_{\ell+1}) < \sigma(\lambda) - \sigma(\alpha_0) + \frac{\varepsilon}{2}. \quad (3.25)$$

Since $\{x_{k_n}\}_{n \geq 1}$ cannot be $T[g]$ -critical (thanks to assumption (iii)), we deduce from Definition 2.12 that it has accumulation points as $n \rightarrow \infty$. We define x_λ to be any accumulation point of $\{x_{k_n}\}_{n \geq 1}$. Then, the last part of the argument is the same as in Case 3 of Initialization: we chose $n_0 \in \mathbb{N}$ in a way that

$$d(x_{k_{n_0}}, x_\lambda) < \frac{\varepsilon}{2T[g](x_{k_0})}$$

and we deduce that

$$\sum_{n=0}^{n_0-1} T[g](x_{k_n}) d(x_{k_n}, x_{k_{n+1}}) + T[g](x_{k_{n_0}}) d(x_{k_{n_0}}, x_\lambda) < \sum_{n=0}^{+\infty} T[g](x_{k_n}) d(x_{k_n}, x_{k_{n+1}}) + \frac{\varepsilon}{2}, \quad (3.26)$$

and the result follows by combining (3.25) and (3.26).

This completes the transfinite induction and allows to obtain a generalized sequence $\{x_\alpha\}_{\alpha < \lambda}$ for all countable ordinal λ . Thus we ultimately define $\{x_\alpha\}_{\alpha < \Omega}$ with $\{g(x_\lambda)\}_{\lambda < \Omega}$ uncountable and strictly decreasing, contradicting that g is bounded from below.

Therefore, $f < (1 + \rho)g$. Repeating the procedure for any $\rho > 0$, we deduce that $f \leq g$ should hold. The proof is complete. \square

Remark 3.4 (discussion on the assumptions). (a). Assumptions (ii) and (iii) are complementary and independent: indeed, asymptotically critical sequences can neither yield nor be obtained by critical points, since they are not allowed to converge (c.f. Definition 3.1). In particular, in absence of compactness, the set of critical points $\mathcal{Z}_T(g)$ can be empty, case in which assumption (ii) of Theorem 3.3 is trivially satisfied and provides no information. This potential lack of information is contemplated by assumption (iii).

(b). The main difficulty in proving Theorem 3.3 is that the construction requires transfinite induction, while assumption (iii) allows comparison through sequences. Indeed, from a point x_α for which $T[g](x_\alpha) > 0$, we produce a descent point $x_{\alpha+}$ (that is, $g(x_{\alpha+}) < g(x_\alpha)$, and after countably many descent points, we select an adequate accumulation point; due to the (possible) existence of points $x \in X \setminus \mathcal{Z}_T(g)$ for which $\liminf_{y \rightarrow x} T[g](y) = 0$, such construction might end prematurely unless we allow to restart the process from the limit points, inducing a transfinite construction (see Example 3.5). The critical part is to prove, using only asymptotically critical sequences, that for every limiting ordinal λ , the accumulation point x_λ can always be constructed. The importance of the invariant properties (P1)–(P3) during this construction was precisely the fact that if such a point x_λ fails to exist, then we would be able to extract an asymptotically critical (cofinal) subsequence from $\{x_\alpha\}_{\alpha < \lambda}$, and compare the functions over that sequence.

(c). Let us momentarily assume that condition (3.3) is imposed to every sequence $\{z_n\}_{n \geq 1} \subset X \setminus \mathcal{Z}_T(g)$ satisfying (3.1), rather than only to those that are free of accumulation points. Let further \bar{z} denote some accumulation point of $\{z_n\}_{n \geq 1}$. Then the case where \bar{z} is critical (i.e. $\bar{z} \in \mathcal{Z}_T(g)$) is already covered by assumption (ii) of Theorem 3.3 (since f is continuous and g

lower semicontinuous) while the case where $\bar{z} \notin \mathcal{Z}_T(g)$ leads to a superfluous assumption making the statement of the theorem weaker. This is illustrated in Example 3.5 below, where the descent operator $T[f]$ is the local slope $s[f]$. The example reveals that accumulation points of sequences satisfying (3.1) might not be critical for the slope $s[g]$ but instead, for the closure of $s[g]$ (called regularized slope in [11]) and that it is neither necessary nor desirable to impose any condition there.

Example 3.5. Set $X = [1, +\infty)$ with the usual distance. For each interval $I_n = [n, n + 1)$ we define the function $g_n : I_n \rightarrow [0, 1]$ given by

$$g_n(x) = \frac{1}{n+1} + \left(\frac{1}{n(n+1)} \right) (x - (n+1))^{n(n+1)}.$$

We finally define $g : [1, +\infty) \rightarrow [0, 1]$ given by $g(x) = g_n(x)$ whenever $x \in I_n$.




Figure 3: Function $g : [1, +\infty) \rightarrow [0, 1]$, constructed by blocks $I_n = [n, n + 1)$. Plot of the first 9 intervals. At each integer point $n \geq 1$, the lateral derivatives are $g'_-(n) = 0$ from the left and $g'_+(n) = 1$ from the right.

Consider the descent operator $T[g] = s[g]$ (local slope) and notice that

$$s[g](x) = (n+1-x)^{n(n+1)-1} > 0, \quad \text{for } x \in [n, n+1) \quad \text{and} \quad \mathcal{Z}_T[g] = \emptyset.$$

Notice further that

$$\inf_{x \in X} g = \lim_{x \rightarrow +\infty} g(x) = 0.$$

Finally, for every $n \in \mathbb{N}$, the point $\bar{x} := n$ is not critical, but it is critical for the regularized slope (see Figure 3), that is:

$$\liminf_{x \rightarrow n} s[g](x) = 0. \tag{3.27}$$

It is easy to see that $f \leq g$ for every continuous function $f : X \rightarrow \mathbb{R}$ satisfying

$$(a) \quad s[f] < s[g] \quad \text{on } X \quad \text{and} \quad (b) \quad \liminf_{x \rightarrow +\infty} f(x) < 0.$$

Indeed, we can either apply Theorem 3.3 or do the following elementary proof: pick any increasing sequence $x_n \rightarrow +\infty$ such that $\lim_{n \rightarrow +\infty} f(x_n) = \liminf_{x \rightarrow +\infty} f(x)$, and assume towards a contradiction that $f(x_0) \geq (1 + \rho)g(x_0)$, for some $\rho > 0$. Following the construction of [7, Lemma 3.3], we can build a (generalized) sequence $\{z_\lambda\}_\lambda$ in the compact interval $[x_0, x_1]$, strictly decreasing for g and such that $f(z_\lambda) \geq (1 + \rho)g(z_\lambda)$ for each λ . The construction eventually ends (due to cardinality obstructions) and the only way for this to happen is that $z_\lambda \rightarrow x_1$, since this is the only critical point of g restricted to $[x_0, x_1]$. Continuity of f and lower semicontinuity of g would yield that $f(x_1) \geq (1 + \rho)g(x_1)$. An inductive argument shows that $f(x_n) \geq (1 + \rho)g(x_n)$ for all $n \in \mathbb{N}$, which yields $\lim_n f(x_n) \geq \inf g = 0$, leading to a contradiction.

On the other hand, if Definition 3.1 allowed to consider convergent sequences, then Theorem 3.3 could not directly apply since it would have required an extra condition on all sequences satisfying (3.27), leading to (infinitely many) unnecessary extra conditions: $f(n) < g(n)$ for all $n \geq 1$.

Finally, observe that for every $n \in \mathbb{N}$, the steepest descent curve $\gamma_n : [0, +\infty) \rightarrow \mathbb{R}$ solving

$$\begin{cases} \dot{\gamma}(t) = -\nabla g(\gamma(t)), & t \geq 0, \\ \gamma(0) = n, \end{cases}$$

satisfies that $\lim_{t \rightarrow +\infty} \gamma_n(t) = n + 1$. Thus, if we follow the construction of Theorem 3.3 by taking a descent point at each iteration, we should obtain a generalized sequence $\{x_\alpha\}$ (similar to a concatenation of discretizations of the curves $\{\gamma_n\}$) diverging to $+\infty$. The delicate construction of the proof of Theorem 3.3 would allow us to retrieve an s -asymptotically critical cofinal subsequence for g , and therefore to compare f and g at the limit values. \diamond

3.2 Determination in complete metric spaces

If two functions f, g have the same descent modulus at every point (that is, $T[f] = T[g]$), then they have the same critical set ($\mathcal{Z}_T := \mathcal{Z}_T(f) = \mathcal{Z}_T(g)$) and the same asymptotically critical sequences ($\mathcal{AZ}_T := \mathcal{AZ}_T(f) = \mathcal{AZ}_T(g)$). Using the same strategy as in Theorem 2.4, we obtain the main result of this work.

Theorem 3.6 (main determination result). *Let T be a metrically compatible descent modulus on the complete metric space (X, d) . Let $f, g \in \bar{\mathcal{C}}(X) \cap \text{dom}(T)$ be bounded from below and satisfy*

$$T[f](x) = T[g](x), \quad \text{for all } x \in X.$$

Assume that $f|_{\mathcal{Z}_T} = g|_{\mathcal{Z}_T}$ and $\liminf_{n \rightarrow \infty} f(z_n) = \liminf_{n \rightarrow \infty} g(z_n)$, for all $\{z_n\}_{n \geq 1} \in \mathcal{AZ}_T$.

Then

$$f(x) = g(x), \quad \text{for all } x \in X.$$

By considering only metric spaces and metrically compatible descent moduli, it is clear that the above result generalizes Theorem 2.4: indeed, for every $f \in \mathcal{F}$ one has that $\mathcal{AZ}_T(f) = \emptyset$ whenever X is compact.

Theorem 3.6 also generalizes [18, Section 4]. This is due to the fact that for the global slope \mathcal{G} defined in (2.13), every \mathcal{G} -asymptotically critical sequence of f is infimizing for the function f . This is the content of the following lemma.

Lemma 3.7 (infimizing sequences). *Let (X, d) be a complete metric space, $f \in \bar{\mathcal{C}}(X) \cap \text{dom}(\mathcal{G})$ and $\{z_n\}_{n \geq 1}$ a \mathcal{G} -asymptotically critical sequence (Definition 3.1). Then*

$$\liminf_{n \rightarrow \infty} f(z_n) = \inf f. \quad (3.28)$$

Proof. Let $\{z_n\}_{n \geq 1}$ be a \mathcal{G} -asymptotically critical sequence for the function f . Then, $\{\mathcal{G}[f](z_n)\}_{n \geq 1}$ is a sequence of strictly positive numbers that converges to zero. We set $k_1 = 1$ and define inductively

$$k_{n+1} := \min \{m \geq k_n : \mathcal{G}[f](z_m) < \mathcal{G}[f](z_{k_n})\}.$$

Then, $\{\mathcal{G}[f](z_{k_n})\}_{n \geq 1}$ is strictly decreasing and for $m \in [k_n, k_{n+1}) \cap \mathbb{N}$ we have

$$\mathcal{G}[f](z_{k_n}) \leq \mathcal{G}[f](z_m).$$

We deduce that for every $n \geq 1$

$$\begin{aligned} \mathcal{G}[f](z_{k_n}) d(z_{k_n}, z_{k_{n+1}}) &\leq \mathcal{G}[f](z_{k_n}) \sum_{k_n \leq m < k_{n+1}} d(z_m, z_{m+1}) \\ &\leq \sum_{k_n \leq m < k_{n+1}} \mathcal{G}[f](z_m) d(z_m, z_{m+1}) \end{aligned}$$

and consequently

$$\sum_{n=1}^{\infty} \mathcal{G}[f](z_{k_n}) d(z_{k_n}, z_{k_{n+1}}) \leq \sum_{m=1}^{\infty} \mathcal{G}[f](z_m) d(z_m, z_{m+1}) < +\infty.$$

Let $u \in X$ be arbitrarily chosen. We deduce from the definition of the global slope (2.13) that

$$f(z_{k_n}) \leq f(u) + \mathcal{G}[f](z_{k_n}) d(z_{k_n}, u), \quad \text{for all } n \geq 1.$$

Therefore, it suffices to show that

$$\liminf_{n \rightarrow \infty} \mathcal{G}[f](z_{k_n}) d(z_{k_n}, u) = 0.$$

To this end, take $n, m \in \mathbb{N}$ with $n < m$. Then, since $\mathcal{G}[f](z_{k_n})$ is decreasing, we deduce:

$$\mathcal{G}[f](z_{k_m}) d(z_{k_m}, u) \leq \mathcal{G}[f](z_{k_{m-1}}) d(z_{k_{m-1}}, z_{k_m}) + \mathcal{G}[f](z_{k_m}) d(z_{k_{m-1}}, u)$$

and consequently

$$\mathcal{G}[f](z_{k_m}) d(z_{k_m}, u) \leq \sum_{n \leq \ell < m} \mathcal{G}[f](z_{k_\ell}) d(z_{k_\ell}, z_{k_{\ell+1}}) + \mathcal{G}[f](z_{k_m}) d(z_{k_n}, u).$$

Thus, keeping n fixed and passing to the limit as $m \rightarrow \infty$, we deduce that:

$$\liminf_{m \rightarrow \infty} \mathcal{G}[f](z_{k_m}) d(z_{k_m}, u) \leq \sum_{\ell=n}^{+\infty} \mathcal{G}[f](z_{k_\ell}) d(z_{k_\ell}, z_{k_{\ell+1}}) \leq \sum_{\ell=n}^{\infty} \mathcal{G}[f](z_\ell) d(z_\ell, z_{\ell+1}).$$

Since the last quantity becomes arbitrarily small as n increases, the conclusion follows. \square

Remark 3.8. Comparing the above lemma with [18, Lemma 4.2], we observe that the main difference is that the latter considers summable sequences where $\sum_n \mathcal{G}[g](z_{n+1})d(z_n, z_{n+1}) < +\infty$. This can be seen as proximal algorithm-type condition. However, in our context, we need to consider asymptotically critical sequences where $\sum_n \mathcal{G}[g](z_n)d(z_n, z_{n+1}) < +\infty$. Thus, asymptotically critical sequences can be seen as gradient algorithm-type condition, which is strongly related on how descent (generalized) sequences are constructed from descent moduli.

Finally, Lemma 3.7, together with the fact that any critical point for the global slope has to be a global minimizer, yields directly the following corollary.

Corollary 3.9 (Global slope determination, [18]). *Let (X, d) be a complete metric space and $f, g : X \rightarrow \mathbb{R}$ be two proper lower semicontinuous functions which are bounded from below and continuous on their domain. If $\mathcal{G}[f](x) = \mathcal{G}[g](x)$, for all $x \in X$ and $\inf f = \inf g$, then $f = g$.*

Corollary 3.10 below reveals that a continuous bounded from below function $f \in \bar{\mathcal{C}}(X) \cap \text{dom}(T)$ necessarily possesses either a critical point or an asymptotically critical sequence.

Corollary 3.10 (existence of critical elements). *Let $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$ be a lower semicontinuous function which is continuous on its domain and bounded from below. If $f \in \text{dom}(T)$ for some metrically compatible descent modulus on X , then either $\mathcal{Z}_T(f) \neq \emptyset$ or $\mathcal{AZ}_T(f) \neq \emptyset$.*

Proof. We may assume that $\inf f = 0$. Fix $\varepsilon > 0$, set $g := (1 + \varepsilon)f$ and $\tilde{f} = f + 1$. It follows that $\inf g = 0 < \inf \tilde{f} = 1$. In view of Proposition 2.16 and property (\mathcal{D}_3) of the descent modulus, we deduce that $\mathcal{Z}_T(g) \subset \mathcal{Z}_T(\tilde{f})$ and $T[g](x) > T[\tilde{f}](x)$, for all $x \in X \setminus \mathcal{Z}_T(g)$. Moreover, if a sequence $\{z_n\}_n$ is $T[g]$ -critical, then it is also $T[\tilde{f}]$ -critical (and consequently, $T[f]$ -critical). Let us assume, towards a contradiction that g has no critical points and no $T[g]$ -critical sequences. Then $\mathcal{Z}_T(g) = \emptyset$ and assumptions (ii) and (iii) of Theorem 3.3 are trivially fulfilled. We deduce that $g > \tilde{f}$ which is a contradiction. Therefore, either $\mathcal{Z}_T(g) \neq \emptyset$ or $\mathcal{AZ}_T(g) \neq \emptyset$. \square

(Open question) It is well-known that the local and the global slopes coincide for convex functions in any Banach space and are equal to the remoteness of the subdifferential (the distance of the convex subdifferential to zero). This fact was used in [18, Section 5] to obtain a nontrivial generalization (from Hilbert to Banach spaces) of the determination result for the class of convex functions obtained in [4, Theorem 3.8] (smooth case) and [17] (nonsmooth case). Indeed, the identification of derivatives with gradients and the use of (sub)gradient systems played a crucial role in the latter works. In the recent work [6], the authors have again used the Hilbertian structure to show that the deviation between the slopes of two convex functions controls the deviation between the functions themselves. It is not known if such slope-based sensitivity result would hold for convex functions in Banach spaces, or more generally, if one can use metric descent modulus deviations to measure deviations of functions in general.

Acknowledgements. The authors wish to thank Sebastian Tapia-García for useful discussions related to Remark 2.8. The first author acknowledges support from the Austrian Science Fund (FWF, P-36344-N). The third author was partially funded by ANID-Chile, through the grant FONDECYT 11220586, and FB210005 BASAL funds for centers of excellence.

References

- [1] L. AMBROSIO, N. GIGLI, G. SAVARÉ, *Gradient flows in metric spaces and in the space of probability measures*, Lectures in Mathematics. Birkhäuser, second edition, 2008.
- [2] D. AZÉ, J.-N. CORVELLEC, Characterizations of error bounds for lower semicontinuous functions on metric spaces, *ESAIM Control Optim. Calc. Var.* **10** (2004), 409–425.
- [3] J.-B. BAILLON, Personal communication (October 2018).
- [4] T. BOULMEZAOUD, P. CIEUTAT, A. DANILIDIS, Gradient flows, second-order gradient systems and convexity, *SIAM J. Optim.* **28** (2018), 2049–2066.
- [5] J. N. CORVELLEC, M. DEGIOVANNI, M. MARZOCCHI. Deformation properties for continuous functionals and critical point theory, *Topol. Methods Nonlinear Anal.* **1** (1993), 151–171.
- [6] A. DANILIDIS, D. DRUSVYATSKIY, The slope robustly determines convex functions, *Proc. Amer. Math. Soc.* **151** (2023), 4751–4756.
- [7] A. DANILIDIS, L. MICLO, D. SALAS, Descent modulus and applications, [arXiv 2211.11819](https://arxiv.org/abs/2211.11819) (preprint 45p, 2022).
- [8] A. DANILIDIS, D. SALAS, A determination theorem in terms of the metric slope, *Proc. Amer. Math. Soc.* **150** (2022), 4325–4333.
- [9] M. DEGIOVANNI, M. MARZOCCHI. A critical point theory for nonsmooth functionals. *Ann. Mat. Pura Appl.* **167** (1994), 73–100.
- [10] E. DE GIORGI, A. MARINO, M. TOSQUES, Problems of evolution in metric spaces and maximal decreasing curve. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat.* **68** (1980), 180–187.
- [11] D. DRUSVYATSKIY, A. D. IOFFE, A. S. LEWIS, Curves of descent, *SIAM J. Control Optim.* **53** (2015), 114–138.
- [12] L. GAJEK, D. ZAGRODNY. Countably orderable sets and their applications in optimization. *Optimization* **26** (1992), 287–301.
- [13] W. GANGBO, A. SWIECH, Metric viscosity solutions of Hamilton-Jacobi equations depending on local slopes, *Calc. Var. Part. Diff. Eq.* **54** (2015), 1183–1218.
- [14] Y. GIGA, N. HAMAMUKI, A. NAKAYASU, Eikonal equations in metric spaces, *Trans. Amer. Math. Soc.* **367** (2015), 49–66.
- [15] Q. LIU, N. SHANMUGALINGAM, X. ZHOU, Equivalence of solutions of eikonal equation in metric spaces, *J. Diff. Eq.* **272** (2021), 979–1014.
- [16] A. D. IOFFE, *Variational analysis of regular mappings*, Springer Monographs in Mathematics. Springer, Cham, 2017.
- [17] P. PEREZ-AROS, D. SALAS, E. VILCHES, Determination of convex functions via subgradients of minimal norm, *Math. Program.* **190** (2021), 561–583.

- [18] L. THIBAUT, D. ZAGRODNY, Determining functions by slopes, *Communications in Contemporary Mathematics* **25** (2023), no 7, 2250014.
- [19] E. VILCHES, Proximal determination of convex functions, *J. Convex Anal.* **28** (2021), 1187–1192.

Aris DANIILIDIS, Tri Minh LE

Institut für Stochastik und Wirtschaftsmathematik, VADOR E105-04
TU Wien, Wiedner Hauptstraße 8, A-1040 Wien

E-mail: {[aris.daniilidis](mailto:aris.daniilidis@tuwien.ac.at), [minh.le](mailto:minh.le@tuwien.ac.at)}@tuwien.ac.at
<https://www.arisdaniilidis.at/>

Research supported by the Austrian Science Fund grant FWF P-36344N.

David SALAS

Instituto de Ciencias de la Ingeniería, Universidad de O'Higgins
Av. Libertador Bernardo O'Higgins 611, Rancagua, Chile

E-mail: david.salas@uoh.cl
<http://davidsalasvidela.cl>

Research supported by the grants:

CMM FB210005 BASAL funds for centers of excellence (ANID-Chile)
FONDECYT 11220586 (Chile)